OPIC



(12) (19) (CA) Demande-Application

C I P O
CANADIAN INTELLECTUAL
PROPERTY OFFICE

(21)(A1) **2,199,636**

(86) 1995/10/05 (87) 1996/04/25

Office de la propriété intellectue*d*e du Canada

Gel. to Carresp. to 310 96/11574

(72) GLOCK, Jutta, CH

(72) HUDETZ, Manfred, US

(72) KERBER, Elmar, DE

(71) NOVARTIS AG, CH

(51) Int.Cl.⁶ A01N 43/56, A01N 25/32, A01N 43/90

(30) 1994/10/17 (3120/94-5) CH

(54) COMPOSITIONS HERBICIDES

(54) HERBICIDAL COMPOSITIONS

(57) Ces compositions herbicides sélectives, destinées à la lutte contre les graminées et les mauvaises herbes dans des cultures de végétaux, comprennent: a) une quantité efficace sur le plan herbicide d'un composé de la formule (I), ainsi que les sels et diastéréoisomères de celui-ci; dans cette formule, R₁ représente le groupe (i), (ii) ou (iii); les substituants R₄ représentent chacun indépendamment halogène, nitro, cyano, alkyle C₁-C₄, haloalkyle C₁-C₄, alcoxy C₁-C₁₀, haloalcoxy C₁-C₄, alcényloxy C₃-C₆, alcoxy C₁-C₄-alcoxy C₂-C₄, C₃-C₆, alkylcarbonyle alcynyloxy alcoxycarbonyle C₁-C₄, alkylthio C₁-C₄, alkylsulfinyle C₁-C₄, alkylsulfonyle C₁-C₄, amino, alkylamino C₁-C₄, ou di-alkylamino C₁-C₄; R₅ représente le groupe (iv) ou (v); n vaut 0, 1, 2, 3 ou 4; m vaut 0 ou 1, la somme de m et de n valant 0, 1, 2, 3 ou 4; q vaut 0, 1, 2 ou 3; X₁ représente oxygène, soufre, -CH2- ou -N(R7)-; les substituants R₆ représentent chacun indépendamment alkyle C₁-C₄, halogène, haloalkyle C₁-C₄, alcoxy C₁-C₄, haloalcoxy C₁-C₄, nitro, cyano, alcoxycarbonyle C₁-C₄, amino, alkylamino C₁-C₄ ou di-alkylamino C₁-C₁; R₇ représente hydrogène, alkyle C₁-C₄, formyle ou alkylcarbonyle C₁-C₄; A et B représentent chacun indépendamment hydrogène, alkyle, alcényle, alcynyle, alcoxyalkyle, alkylthioalkyle ou cycloalkyle, ou aryle substitué ou non substitué; ou bien A et B pris ensemble forment le radical divalent d'un système mono, bi, tri ou polycyclique, saturé ou non, substitué ou non; G représente hydrogène ou un groupe (a) -CO-R₁₈, (b), (c) -SO₂-R₂₀, (d), (e) ou (f) X; L et M représentent chacun indépendamment oxygène ou soufre; R₁₈ représente (57) Selective herbicidal compositions for controlling grasses and weeds in crops of cultivated plants, comprising: a) a herbicidally effective amount of a compound of formula (I), wherein R₁ is the (i), (ii) or (iii) group; the substituents R₄ are each independently of one another halogen, nitro, cyano, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₁₀alkoxy, C₁-C₄haloalkoxy, C₃- C_6 alkenyloxy, C_1 - C_4 alkoxy- C_2 - C_4 alkoxy, C₁-C₁alkylcarbonyl, C₆alkynyloxy, C₁alkoxycarbonyl, C₁-C₁alkylthio, C₁-C₂alkylsulfinyl, C_1 - C_4 alkylsulfonyl, amino, C_1 - C_4 alkylamino or di- C_1 - C_A alkylamino; R_5 is the (iv) or (v) group; n is 0, 1, 2, 3 or 4; m is 0 or 1, the sum of m and n being 0, 1, 2, 3 or 4; q is 0, 1, 2 or 3; X_1 is oxygen, sulfur, -CH₂- or -N(R₇)-; the substituents R₆ are each independently of one another C_1 - C_4 alkyl, halogen, C_1 - C_4 haloalkyl, C_1 -C₁alkoxy, C₁-C₂haloalkoxy, nitro, cyano, C₁-C₁alkoxycarbonyl, amino, C₁-C₂alkylamino or di-C₁-C₄alkylamino; R₇ is hydrogen, C₁-C₄alkyl, formyl or C₁-C₄alkylcarbonyl; A and B are each independently of one another hydrogen, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkylthioalkyl or cycloalkyl, or substituted or unsubstituted aryl; or A and B, taken together, form the divalent radical of a saturated or unsaturated and unsubstituted or substituted mono-, bi-, tri- or polycyclic system; G is hydrogen or a group -CO-R₁₈ (a), (b), - SO_2 - R_{20} (c), (d), (e) or X (f); L and M are each independently of the other oxygen or sulfur; R₁₈ is halogen-substituted alkyl, alkenyl, alkoxyalkyl, alkylthioalkyl, polyalkoxyalkyl or cycloalkyl, which may contain hetero atoms; unsubstituted or substituted phenyl, unsubstituted or substituted phenylalkyl, alkyle,

THIS PAGE BLANK (US.O.T.)



(21) (A1) **2,199,636** (86) 1995/10/05

(87) 1995/10/05 (87) 1996/04/25

alcoxyalkyle, alkylthioalkyle, alcényle, polyalcoxyalkyle ou cycloalkyle substitués par halogène et pouvant contenir des hétéroatomes; phényle non substitué ou substitué, phénylalkyle non substitué ou phénoxyalkyle hétéroaryle substitué, substitué; R₁₉ substitué, ou hétéroaryloxyalkyle alkyle, alcénvle. alcoxyalkyle représente polyalcoxyalkyle substitués par halogène, ou phényle ou benzyle non substitués ou substitués; R₂₀, R₂₁ et R₂₂ représentent chacun indépendamment alkyle, alcoxy, dialkylamino, alkylthio, alcénylthio, alkylamino, alcynylthio ou cycloalkylthio non substitués ou substitués par halogène, ou bien phényle, phénoxy ou phénylthio non substitués ou substitués; R₂₃ et R₂₄ représentent chacun indépendamment hydrogène, alkyle, alcényle, alcoxy ou alcoxyalkyle non substitués ou substitués par halogène, phényle ou benzyle non substitués ou substitués; ou bien R₂₃ et R₂₄, pris ensemble, forment un radical alcényle qui peut contenir oxygène en tant qu'hétéroatome; et X représente un équivalent d'un ion métal ou un ion ammonium; et b) afin d'exercer une action antagoniste à l'égard de cet herbicide, une quantité efficace en tant qu'antidote de: soit un dérivé quinoléine de la formule (IIa) dans laquelle R_{10} représente hydrogène, alkyle C_1 - C_8 ou alkyle C_1 -C₈ substitué par alcoxy C₁-C₆ ou alcényloxy C₃-C₆; et X₂ représente hydrogène ou chloro, soit un dérivé de l'acide 1-phénylazole-3-carboxylique de la formule (IIb) dans laquelle E représente azote ou méthine; R₁₁ représente -CCl3 ou phényle non substitué ou substitué par halogène; R₁₂ et R₁₃ représentent chacun indépendamment hydrogène ou halogène; et R₁₄ représente alkyle C₁-C₄. Ces composés sont particulièrement appropriés à la lutte contre les mauvaises herbes dans des cultures de végétaux, notamment le maïs et les céréales.

substituted heteroaryl, substituted phenoxyalkyl, or substituted heteroaryloxyalkyl; R₁₀ is halogensubstituted alkyl, alkenyl, alkoxyalkyl polyalkoxyalkyl, or unsubstituted or substituted phenyl or benzyl; R_{20} , R_{21} and R_{22} are each independently of one another unsubstituted or halogen-substituted alkyl, alkoxy, alkylamino, dialkylamino, alkynylthio cycloalkylthio, alkenylthio, or unsubstituted or substituted phenyl, phenoxy or phenylthio; R₂₃ and R₂₄ are each independently of the other hydrogen, unsubstituted or halogen-substituted alkyl, alkenyl, alkoxy or alkoxyalkyl, unsubstituted or substituted phenyl or benzyl; or R23 and R24, taken together, form an alkenyl radical, which may contain oxygen as hetero atom; and X is a metal ion equivalent or an ammonium ion; as well as salts and diastereoisomers of the compounds of formula (I); and b) to antagonise the herbicide, an antidotally effective amount of either a quinoline derivative of formula (IIa), wherein R₁₀ is hydrogen, C₁-C₈alkyl, or C₁-C₈alkyl which is substituted by C₁-C₆alkoxy or C₃-C6alkenyloxy; and X2 is hydrogen or chloro; of a 1phenylazole-3-carboxylic acid derivative of formula (IIb), wherein E is nitrogen or methine; R₁₁ is -CCl₃ or unsubstituted or halogen-substituted phenyl; R₁₂ and R₁₃ are each independently of the other hydrogen or halogen; and R₁₄ is C₁-C₄alkyl, are particularly suitable for controlling weeds in crops of cultivated plants, in particular maize and cereals.

THIS PAGE BLANK (USOTO)

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶:

A1

(11) International Publication Number:

WO 96/11574

A01N 43/90, 43/56, 25/32

(43) International Publication Date:

25 April 1996 (25.04.96)

(21) International Application Number:

PCT/EP95/03935

(22) International Filing Date:

5 October 1995 (05.10.95)

(30) Priority Data:

3120/94-5

17 October 1994 (17.10.94) CH

(71) Applicant (for all designated States except US): CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): GLOCK, Jutta [DE/CH]; Rifeldweg 2, CH-4322 Mumpf (CH). HUDETZ, Manfred [DE/US]; 1302D Adams Farm Parkway, Greensboro, NC 27407 (US). KERBER, Elmar [DE/DE]; Niederwihl 131, D-79733 Görwihl (DE).
- (74) Common Representative: CIBA-GEIGY AG; Patentabteilung, Klybeckstrasse 141, CH-4002 Basle (CH).

(81) Designated States: AL, AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).

Published

With international search report.

2199636

(54) Title: HERBICIDAL COMPOSITIONS

(57) Abstract

Selective herbicidal compositions for controlling grasses and weeds in crops of cultivated plants, comprising: a) a herbicidally effective amount of a compound of formula (I), wherein R₁ is the (i), (ii) or (iii) group; the substituents R4 are each independently of one another halogen, nitro, cyano, C1-C4alkyl, C1-C4haloalkyl, C1-C10alkoxy, C1-C4haloalkoxy, C3-C6alkenyloxy, C1-C4alkoxy-C2-C4alkoxy, C3-C6alkynyloxy, C1-C4alkylcarbonyl, C1-C4alkoxycarbonyl, C_1 -Calkylthio, C_1 -Calkylsulfinyl, C_1 -Calkylsulfonyl, amino, C_1 -Calkylamino or di- C_1 -Calkylamino; R_5 is the (iv) or (v) group; n is 0, 1, 2, 3 or 4; m is 0 or 1, the sum of m and n being 0, 1, 2, 3 or 4; q is 0, 1, 2 or 3; X_1 is oxygen, sulfur, -CH₂- or -N(R_7)-; the substituents R6 are each independently of one another C1-C4alkyl, halogen, C1-C4haloalkyl, C1-C4alkoxy, C1-C4haloalkoxy, nitro, cyano, C1-C4alkoxycarbonyl, amino, C1-C4alkylamino or di-C₁-C₄alkylamino; R₇ is hydrogen, C₁-C₄alkyl, formyl or C₁-C₄alkylcarbonyl; A and B are each independently of one another hydrogen, alkyl, alkenyl, alkoxyalkyl, alkylthioalkyl or cycloalkyl, or substituted or unsubstituted aryl; or A and B, taken together, form the divalent radical of a saturated or unsaturated and unsubstituted or substituted mono-, bi-, trior polycyclic system; G is hydrogen or a group -CO-R₁₈ (a), (b), -SO₂-R₂₀ (c), (d), (e) or X (f); L and M are each independently of the other oxygen or sulfur; R₁₈ is halogen-substituted alkyl, alkenyl, alkoxyalkyl, alkylthioalkyl, polyalkoxyalkyl or cycloalkyl, which may contain hetero atoms; unsubstituted or substituted phenyl, unsubstituted or substituted phenylalkyl, substituted heteroaryl, substituted phenoxyalkyl, or substituted heteroaryloxyalkyl; R19 is

halogen-substituted alkyl, alkenyl, alkoxyalkyl or polyalkoxyalkyl, or unsubstituted or substituted phenyl or benzyl; R₂₀, R₂₁ and R₂₂ are each independently of one another unsubstituted or halogen-substituted alkyl, alkoxy, alkylamino, dialkylamino, alkylthio, alkenylthio, alkynylthio or cycloalkylthio, or unsubstituted or substituted phenyl, phenoxy or phenylthio; R₂₃ and R₂₄ are each independently of the other hydrogen, unsubstituted or halogen-substituted alkyl, alkenyl, alkoxy or alkoxyalkyl, unsubstituted or substituted phenyl or benzyl; or R₂₃ and R₂₄, taken together, form an alkenyl radical, which may contain oxygen as hetero atom; and X is a metal ion equivalent or an ammonium ion; as well as salts and diastereoisomers of the compounds of formula (I); and b) to antagonise the herbicide, an antidotally effective amount of either a quinoline derivative of formula (IIa), wherein R₁₀ is hydrogen, C₁-C₅alkyl, or C₁-C₅alkyl which is substituted by C₁-C₅alkoxy or C₃-C₅alkenyloxy; and X₂ is hydrogen or chloro; of a 1-phenylazole-3-carboxylic acid derivative of formula (IIb), wherein E is nitrogen or methine; R₁₁ is -CCl₃ or unsubstituted or halogen-substituted phenyl; R₁₂ and R₁₃ are each independently of the other hydrogen or halogen; and R₁₄ is C₁-C₅alkyl, are particularly suitable for controlling weeds in crops of cultivated plants, in particular maize and cereals.

THIS PAGE BLANK (US.07.9)

Herbicidal compositions

The present invention relates to a selective herbicidal composition for controlling grasses and weeds in crops of cultivated plants, especially in crops of maize and cereals, which composition comprises a herbicide and a safener (antidote) and protects the cultivated plants, but not the weeds, from the phytotoxic action of the herbicide, and to the use of said composition for controlling weeds in crops of cultivated plants.

When applying herbicides, the cultivated plants may also suffer severe damage owing to factors that include the concentration of the herbicide and the mode of application, the cultivated plant itself, the nature of the soil, and the climatic conditions such as exposure to light, temperature and rainfall.

To counteract this problem and similar ones, the proposal has already been made to use different compounds as safeners which are able to antagonise the harmful action of the herbicide on the cultivated plant, i.e. to protect the cultivated plant while leaving the herbicidal action on the weeds to be controlled virtually unimpaired. It has, however, been found that the proposed safeners often have a very specific action, not only with respect to the cultivated plants but also to the herbicide, and in some cases also subject to the mode of application, i.e. a specific safener will often be suitable only for a specific cultivated plant and a specific class of herbicide or a specific herbicide. Thus, for example, EP-A-0 094 349 discloses quinoline derivatives that protect cultivated plants from the phytotoxic action of herbicides of specific substance classes including chloroacetanilides, phenoxypropioniate herbicides, ureas, triazines, carbamates or diphenyl ethers. EP-A-0 558 448 discloses 1,5-diphenylpyrazole-3-carboxylic acid derivatives for protecting cultivated plants from the phytotoxic action of sulfonyl ureas.

It has now been found that the safeners selected from the two compound classes of the quinoline derivatives and 1-phenylazole-3-carboxylic acid derivatives are suitable for protecting cultivated plants from the phytotoxic action of 3-hydroxy-4-aryl-5-oxopyrazoline derivatives.

Accordingly, the invention provides a selective herbicidal composition comprising, in addition to customary inert formulation assistants such as carriers, solvents and wetting agents, a mixture of

a) a herbicidally effective amount of a herbicide of formula I

$$O - G$$
 R_1
 R_1
 R_1
 R_1
 R_1
 R_1
 R_1
 R_1

wherein

$$R_1$$
 is the $(R_4)_n$, $(R_5)_m$ or $(R_4)_n$ group;

the substituents R_4 are each independently of one another halogen, nitro, cyano, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_{10} alkoxy, C_1 - C_4 haloalkoxy, C_3 - C_6 alkenyloxy, C_1 - C_4 alkoxy- C_2 - C_4 alkoxy, C_3 - C_6 alkynyloxy, C_1 - C_4 alkylcarbonyl, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, amino, C_1 - C_4 alkylamino or di- C_1 - C_4 alkylamino;

$$R_5$$
 is the $-x_1$ or $-x_1$ q group;

n is 0, 1, 2, 3 or 4;

m is 0 or 1, the sum of m and n being 0, 1, 2, 3 or 4;

q is 0, 1, 2 or 3;

 X_1 is oxygen, sulfur, -CH₂- or -N(R₇)-;

the substituents R_6 are each independently of one another C_1 - C_4 alkyl, halogen, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, nitro, cyano, C_1 - C_4 alkoxycarbonyl, amino, C_1 - C_4 alkylamino or di- C_1 - C_4 alkylamino;

R₇ is hydrogen, C₁-C₄alkyl, formyl or C₁-C₄alkylcarbonyl;

A and B are each independently of the other hydrogen, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkylthioalkyl or cycloalkyl, or substituted or unsubstituted aryl; or A and B, taken together, form the divalent radical of a saturated or unsaturated and unsubstituted or substituted mono-, bi-, tri- or polycyclic system;

G is hydrogen or a group -CO-R₁₈ (a),
$$M-R_{19}$$
 (b), -SO₂-R₂₀ (c), R_{21} (d),

$$R_{24}$$
 (e) or X (f);

L and M are each independently of the other oxygen or sulfur;

R₁₈ is halogen-substituted alkyl, alkenyl, alkoxyalkyl, alkylthioalkyl, polyalkoxyalkyl or cycloalkyl, which may contain hetero atoms; unsubstituted or substituted phenyl, unsubstituted or substituted phenylalkyl, substituted heteroaryl, substituted phenoxyalkyl, or substituted heteroaryloxyalkyl;

R₁₉ is halogen-substituted alkyl, alkenyl, alkoxyalkyl or polyalkoxyalkyl, or unsubstituted or substituted phenyl or benzyl;

 R_{20} , R_{21} and R_{22} are each independently of one another unsubstituted or halogen-substituted alkyl, alkoxy, alkylamino, dialkylamino, alkylthio, alkenylthio, alkynylthio or cycloalkylthio, or unsubstituted or substituted phenyl, phenoxy or phenylthio;

 R_{23} and R_{24} are each independently of the other hydrogen, unsubstituted or halogen-substituted alkyl, alkenyl, alkoxy or alkoxyalkyl, unsubstituted or substituted phenyl or benzyl; or

 R_{23} and R_{24} , taken together, form an alkenyl radical, which may contain oxygen as hetero atom; and

X is a metal ion equivalent or an ammonium ion; as well as salts and diastereoisomers of the compounds of formula I; and

b) to antagonise the herbicide, an antidotally effective amount of either a quinoline derivative of formula IIa

- 4 -

wherein

 R_{10} is hydrogen, C_1 - C_8 alkyl, or C_1 - C_8 alkyl which is substituted by C_1 - C_6 alkoxy or C_3 - C_6 alkenyloxy; and X_2 is hydrogen or chloro; of a 1-phenylazole-3-carboxylic acid derivative of formula IIb

$$R_{11}$$
 R_{12}
 R_{13}
(IIb),

wherein

E is nitrogen or methine;

 R_{11} is -CCl₃ or unsubstituted or halogen-substituted phenyl; R_{12} and R_{13} are each independently of the other hydrogen or halogen; and R_{14} is C_1 -C₄alkyl; or of a compound of formula IIb₁

$$R_{67}OOC$$
 R_{68}
 R_{12}
 R_{13}
(IIb₁),

wherein R_{12} and R_{13} have the meanings given above, and R_{66} , R_{67} and R_{68} are each independently of one another C_1 - C_4 alkyl.

The alkyl, alkenyl and alkynyl groups contain, unless specifically mentioned, preferably 1 or 2 to 8 carbon atoms and may be straight-chain or branched, as also applies to the alkyl, alkenyl and alkynyl moiety of the haloalkyl, alkoxy, haloalkoxy, alkoxyalkyl, alkoxyalkoxy, alkenyloxy, alkynyloxy, alkylcarbonyl, alkoxycarbonyl, alkylthio,

alkenylthio, alkynylthio, alkylthioalkyl, alkylsulfinyl, alkylsulfonyl, alkylamino and dialkylamino groups.

The alkyl groups occurring in the substituent definitions are typically methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl or tert-butyl, as well as the isomeric pentyls, hexyls, heptyls and octyls. The alkyl groups by themselves or as moieties of other substituents preferably contain 1 to 8 carbon atoms.

In the above definitions halogen will be taken to mean fluoro, chloro, bromo and iodo, preferably fluoro, chloro and bromo.

Alkenyl will be typically vinyl, allyl, methallyl, 1-methylvinyl or but-2-en-1-yl. Alkenyl radicals containing 2 to 8 carbon atoms in the chain are preferred. The alkenyl radicals are preferably linked via a saturated carbon atom to a hetero atom.

Alkynyl will typically be propargyl, but-2-yn-1-yl, 2-methylbutyn-2-yl, but-3-yn-2-yl and pent-4-yn-1-yl. Alkynyl radicals containing 2 to 8 carbon atoms in the chain are preferred. The alkynyl radicals are preferably linked via a saturated carbon atom to a hetero atom.

Haloalkyl radicals preferably contain 1 to 8 carbon atoms in the chain. Haloalkyl is typically fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 2-fluoroethyl, 2-chloroethyl and 2,2,2-trichloroethyl. Trichloromethyl, difluoromethyl, difluoromethyl, trifluoromethyl and dichlorofluoromethyl are preferred.

Alkoxy radicals preferably contain 1 to 6 carbon atoms in the chain. Alkoxy is typically methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy and tert-butoxy as well as the pentoxy and hexyloxy isomers. Methoxy and ethoxy are preferred.

Alkylcarbonyl is preferably acetyl and propionyl.

Alkoxycarbonyl is methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, n-butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl and tert-butoxycarbonyl. Methoxycarbonyl and ethoxycarbonyl are preferred.

Haloalkoxy radicals preferably contain 1 to 8 carbon atoms in the chain. Haloalkoxy is typically fluoromethoxy, difluoromethoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy,

1,1,2,2-tetrafluoroethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2,2-difluoroethoxy and 2,2,2-trichloroethoxy. Difluoromethoxy, 2-chloroethoxy and trifluoromethoxy are preferred.

Alkylthio radicals preferably contain 1 to 8 carbon atoms in the chain. Alkylthio is typically methylthio, ethylthio, propylthio, isopropylthio, n-butylthio, isobutylthio, sec-butylthio or tert-butylthio. Mmethylthio and ethylthio are preferred.

Alkylsulfinyl is methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl, n-butylsulfinyl, isobutylsulfinyl, sec-butylsulfinyl, tert-butylsulfinyl. Methylsulfinyl and ethylsulfinyl are preferred.

Alkylsulfonyl is methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, isobutylsulfonyl, sec-butylsulfonyl, tert-butylsulfonyl. Methylsulfonyl and ethylsulfonyl are preferred.

Alkoxyalkoxy radicals preferably contain 1 to 8 carbon atoms in the chain. Illustrative examples of alkoxyalkoxy are methoxymethoxy, methoxyethoxy, methoxypropoxy, ethoxymethoxy, ethoxyethoxy, propoxymethoxy and butoxybutoxy.

Alkylamino is typically methylamino, ethylamino, n-propylamino, isopropylamino and the isomeric butylamines.

Dialkylamino is typically dimethylamino, methylethylamino, diethylamino, n-propylmethylamino, dibutylamino and di-isopropylamino. Alkylamino radicals containing 1 to 8 carbon atoms in the chain are preferred.

Alkoxyalkyl radicals preferably contain 1 to 8 carbon atoms. Alkoxyalkyl is typically methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, n-propoxymethyl, n-propoxymethyl, isopropoxymethyl and isopropoxyethyl.

Alkylthioalkyl radicals preferably contain 1 to 8 carbon atoms. Alkylthioalkyl is typically methylthiomethyl, methylthioethyl, ethylthiomethyl, ethylthioethyl, n-propylthiomethyl, n-propylthiomethyl, isopropylthioethyl, butylthiomethyl, butylthiomethyl, butylthioethyl or butylthiobutyl.

Cycloalkyl radicals preferably contain 3 to 8 ring carbon atoms, and are typically cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. These cycloalkyl radicals may be interrupted by hetero atoms such as oxygen atoms and/or sulfur atoms and/or nitrogen atoms.

Phenyl, also as moiety of a substituent such as phenoxy, phenylthio, phenylalkyl, phenoxyalkyl, can generally be unsubstituted or substituted. The substituents may then be in ortho-, meta- and/or para-position. Preferred substituent positions are the ortho- and para-position to the ring linkage site. Preferred substituents are halogen, nitro, cyano, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 alkylthio, C_1 - C_4 haloalkyl.

A and B defined as aryl are α -naphthyl or β -naphthyl, preferably phenyl, and these aromatic rings may carry one or more than one identical or different substituent, typically halogen, nitro, cyano, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 alkylthio, C_1 - C_4 haloalkylthio, C_1 - C_4 haloalkyl.

Heteroaryl in the definition of the radical R_{18} is preferably a five- or six-membered aromatic heterocyclic ring, typically 2-, 3- or 4-pyridyl, pyrimidyl, pyrazinyl, furanyl, thienyl, oxazolyl or isoxazolyl.

Illustrative examples of saturated or unsaturated and unsubstituted or substituted mono-, bi-, tri- or polycyclic systems which the substituents A and B, together with the two linking nitrogen atoms of the pyrazoline ring may form, are:

suitable substituents being halogen, unsubstituted or halogen-substituted C_1 - C_{16} alkyl, C_2 - C_{16} alkenyl, C_1 - C_1 6alkoxy- C_2 - C_6 alkyl, C_1 - C_6 polyalkoxy- C_2 - C_6 alkyl, or phenyl or benzyl which are unsubstituted or substituted by halogen, nitro, C_1 - C_4 alkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkyl; and C_1 0 is halogen, or unsubstituted or halogen-substituted C_1 alkoxy or C_2 alkoxy.

Equivalent meanings may also be obtained for the substituents of composite radicals, typically alkoxycarbonyl, alkylcarbonyl, alkylsulfinyl, alkylsulfonyl, alkenyloxy, alkynyloxy, alkenylthio, alkynylthio, cycloalkylthio and heteroaryloxyalkyl.

By reason of their chemical constitution, the compounds of formula I, wherein G is hydrogen, can be obtained in the tautomeric equilibrium forms Ib \rightleftharpoons Ic \rightleftharpoons Id:

$$\begin{array}{c} A \\ N \\ B \end{array}$$

$$\begin{array}{c} A \\ N \\ O \\ \end{array}$$

$$\begin{array}{c} A \\ N \\ \end{array}$$

The invention also embraces the salts which the compounds of formula I may form together with acids. Suitable salts for the formation of the acid addition salts are organic as well as inorganic acids. Examples of such acids are, inter alia, hydrochloric acid, hydrobromic acid, nitric acid, various phosphoric acids, sulfuric acid, acetic acid, propionic acid, butyric acid, valeric acid, oxalic acid, malonic acid, fumaric acid, lactic acid, tartaric acid and salicylic acid.

Furthermore, specific substituents R₁ to R₇, R₁₈ to R₂₄, A and B, singly or in conjunction

with each other or in conjunction with the basic structure to which they are linked, may have chirality. The invention embraces the racemate as well as the enriched and optically pure forms of the respective stereoisomers.

In the processes disclosed in this specification, the asymmetrically substituted compounds of formula I are usually obtained as racemates, unless chiral educts are used. The stereoisomers can then be isolated in accordance with per se known methods, such as fractionated crystallisation after salt formation with optically pure bases, acids or metal complexes, or chromatographic processes based on physicochemical properties.

The compounds of formula I, wherein A and B are alkyl, alkenyl or alkynyl radicals, are derivatives of the pyrazolidine-3,5-dione system. Where A and B are defined as a saturated or partially unsaturated C₄carbon bridge, formula I is based on the ring system of the 1H-pyrazolo[1,2-a]pyridazine, and, where R₂ and R₃ are defined as a saturated or partially unsaturated C₃carbon bridge, formula I is based on the ring system of the 1H,5H-pyrazolo[1,2-a]pyrazole. The individual ring positions are numbered in accordance with Chemical Abstracts:

1H-pyrazolo[1,2-a]pyridazine

1H,5H-pyrazolo[1,2-a]pyrazole

Compounds of formula I which are preferred for use in the novel composition are those wherein R_1 is the $\binom{(R_4)_n}{(R_5)_m}$ group. Among these compounds, those compounds

are particularly preferred wherein R_4 is C_1 - C_4 alkyl; n is 1, 2, 3 or 4; and m is 0.

Among these compounds, those compounds are also particularly preferred wherein R_5 is the $-x_1$ group.

Particularly important compositions are those wherein R_4 is C_1 - C_4 alkyl, trifluoromethyl, fluoro, chloro, bromo, C_1 - C_{10} alkoxy, allyloxy, propargyloxy, difluoromethoxy,

trifluoromethoxy, methoxyethylenoxy, cyano, nitro, di- C_1 alkylamino or di- C_2 alkylamino, acetyl, C_1 - C_3 alkoxycarbonyl, methylsulfonyl, methylsulfinyl or methylmercapto; R_5 is the

$$-x_1$$
 or $-x_1$ $(R_6)_q$ group; X_1 is oxygen, sulfur, -CH₂-

or -N(CHO)-; R_6 is fluoro, chloro or trifluoromethyl; n is 0, 1, 2 or 3; m is 0 or 1; and 0 is 0, 1 or 2, the sum of m and n being 0, 1, 2 or 3.

Furthermore, those compositions are particularly preferred, wherein R₁ is the

$$(R_4)_n$$
 group; R_4 is fluoro, chloro, methyl, nitro, trifluoromethyl or

methylsulfonyl; n is 0, 1 or 2; and m is 0.

Particularly important compositions are also those, wherein R₁ is the

$$(R_4)_n$$
 group; R_4 is methyl, methoxy, fluoro, chloro or nitro; and n is 0 or 1.

Also preferred are those compositions comprising a safener of formula IIc

Likewise preferred are compositions comprising a safener of formula IId

- 11 -

Further preferred compositions are those comprising a safener of formula He

The following non-limitative Examples illustrate the invention in more detail.

Example P1: Preparation of 2-(2,4,6-trimethylphenylacetyl)-1-ethoxycarbonyl hexahydropyridazine

With stirring, 13.8 g (70 mmol) of mesitylene acetyl chloride in 100 ml of diethyl ether are added dropwise at 20-25°C to a solution of 11.0 g (70 mmol) of 1-ethoxycarbonyl hexahydropyridazine and 10.8 ml (70 mmol) of triethylamine in 350 ml of diethyl ether. Stirring is continued for a further 3 hours at room temperature. The precipitated triethylamine hydrochloride is then removed by suction filtration, and the filtrate is concentrated under vacuum and the residue is chromatographed with ethyl acetate/hexane (1:1) over silica gel, giving 20.1 g (90,5 %) of the desired title compound.

- 12 -

Example P2: Preparation of 2-(2,4,6-trimethylphenyl)-5,6,7,8-tetrahydro-1H-pyrazolo[1,2-a]pyridazine-1,3(2H)dione

$$CH_3$$
 CH_3
 CH_3
 CH_3
 CH_3
 CH_3

3.69 g (88 mmol) of a 60 % suspension of sodium hydride in white oil are added to 75 ml of toluene. To this mixture are then added dropwise 22.3 g (70 mmol) of a solution of 2-(2,4,6-trimethylphenylacetyl)-1-ethoxycarbonyl hexahydropyridazine (Example P1) in 75 ml of toluene at room temperature and the mixture is heated for 6 hours to the boil. While cooling with ice, 10 ml of ethanol are then added dropwise, the reaction mixture is concentrated to dryness under vacuum and the residue is dissolved in 200 ml of 1N NaOH. The product is precipitated from the resultant solution by addition of concentrated hydrochloric acid at O°C. The crude product is purified by recrystallisation from methanol to give 8.9 g of crystals of the desired title product, m.p. 244 - 246°C.

The compounds of Tables 1 to 15 and 19 can be prepared in general accordance with the above Examples and the procedures described therein.

Preferred compounds of formula I are listed in the following Tables 1-15 and 19 as compounds of formula Ia and Ie to Is.

- 13 -

Table 1: Compounds of formula Ia

$$\bigcap_{i}^{N} \bigcap_{O}^{(R_4)_n} (R_5)_m$$
 (Ia)

Cmpd	R ₄	R ₄	R_4	R ₅	
1.001		H			
1.002		2-CH ₃			
1.003		4-CH ₃			
1.004		2-CH ₃	4-CH ₃		
1.005		2-CH ₃	6-CH ₃		
1.006		2-CH ₃	5-CH ₃		
1.007		3-CH ₃	5-CH ₃		
1.008		2-CH ₃	3-CH ₃		
1.009		3-CH ₃	4-CH ₃		
1.010	2-CH ₃	4-CH ₃	6-CH ₃		
1.011	2-CH ₃	4-CH ₃	5-CH ₃		
1.012		2-Cl			
1.013		4-C1			
1.014		2-CI	4-Cl		
1.015		2-Cl	6-C1		
1.016		2-C1	6-F		
1.017		2-CH ₃	4-C1		
1.018		2-CH ₃	4-F		
1.019		2-CI	4-CH ₃		
1.020		2-C1	6-CH ₃		
1.021		2-F	4-F		
1.022		2-F	6-F		
1.023		2-CH ₃	4-O-CH ₃		
1.024		2-CH ₃	6-O-CH ₃		
1.025		2-C1	4-O-CH ₃		
1.026		2-Cl	6-O-CH ₃		

- 14 -

Cmpd R ₄	R ₄	R ₄	R ₅
1.027	3-OCH ₃	4-OCH ₃	
1.028	2-OCH ₃	5-OCH ₃	
1.029	2-OCH ₃	4-OCH ₃	
1.030	2-OCH ₃	6-OCH ₃	
1.031	2-CF ₃	6-CF ₃	
1.032	2-CF ₃	4-CF ₃	
1.033	3-CF ₃	5-CF ₃	
1.034	2-Cl	4-CF ₃	
1.035	2-Cl	6-CF ₃	
1.036	2-NO ₂	4-NO ₂	
1.037	2-Cl	4-NO ₂	
1.038	2-CH ₃	4-NO ₂	
1.039	2-O-CH ₃	4-NO ₂	
1.040	2-F	6-NO ₂	
1.041	2-Cl	6-NO ₂	
1.042	2-CH ₃	6-NO ₂	
1.043	2-O-CH ₃	6-NO ₂	
1.044	2-F	4-NO ₂	
1.045	2-CH ₃	$4-N(C_2H_5)_2$	
1.046	2-Cl	4-SO ₂ -CH ₃	
1.047	2-Cl	4-SO-CH ₃	
1.048	2-Cl	4-S-CH ₃	
1.049	2-Cl	6-SO ₂ -CH ₃	
1.050	2-C1	6-SO-CH ₃	
1.051	2-Cl	6-S-CH ₃	
1.052	2-CH ₃	4-SO ₂ -CH ₃	
1.053	2-CH ₃	4-SO-CH ₃	
1.054	2-CH ₃	4-S-CH ₃	
1.055	2-CH ₃	6-SO ₂ -CH ₃	
1.056	2-CH ₃	6-SO-CH ₃	
1.057	2 -CH $_3$	6-S-CH ₃	
1.058	2-O-CH ₃	6-SO ₂ -CH ₃	
1.059	2-O-CH ₃	6-SO-CH ₃	
1.060	2-O-CH ₃	6-S-CH ₃	
1.061	2-O-CH ₃	4-SO ₂ -CH ₃	
1.062	2-O-CH ₃	4-SO-CH ₃	

- 15 -

Cmpd	R ₄	R ₄	R ₄	R ₅	
1.063		2-O-CH ₃	4-S-CH ₃		
1.064		2-CH ₃	$6-N(C_2H_5)_2$		
1.065		2-C1	6-N(CH ₃) ₂		
1.066		2-Cl	4-N(CH ₃) ₂		
1.067		2-Cl	4-CO ₂ CH ₃		
1.068		2-CH ₃	6-CO ₂ C ₂ H ₅		
1.069		2-CH ₃	$4-CO_2C_2H_5$		•
1.070		2-CH ₃	4-CN		
1.071		2-CH ₃	6-CN		
1.072		2-C1	4-CN		
1.073		2-C1	6-CN		
1.074		2-Cl	4-CO-CH ₃		
1.075		2-O-CHF ₂	4-O-CHF ₂		
1.076		2-CH ₃	4-O-CHF ₂		
1.077		2-Cl	4-O-CF ₃		
1.078		2-O-CF ₃	4-O-CH ₃	•	
1.079		2-O-CHF ₂	4-C1		
1.080		2-O-CHF ₂	6-CH ₃		÷
1.081		2-O-CHF ₂	6-C1		
1.082	2-O-CHF ₂	4-CH ₃	6-CH ₃		
1.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃		
1.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇		
1.085	2-CH ₃	4-O-CH ₃	6-CH ₃		
1.086	2-C1	4-CF ₃	6-C1		
1.087	2-Cl	4-CF ₃	6-F		
1.088	2-C1	4-NO ₂	6-C1		
1.089	2-Cl	4-Cl	6-C1		
1.090	2-F	4-F	6-F		
1.091	2-CH ₃	4-NO ₂	6-CH ₃		
1.092	2-Cl	4-Cl	6-CH ₃		
1.093	2-C1	4-O-CH ₃	6-C1		
1.094	2-Cl	4-Cl	6-O-CH ₃		
1.095	2-F	4-O-CH ₃	6-F		
1.096	2-O-CH ₃	4-CH ₃	6-O-CH ₃		
1.097	•	-	6-CH ₃		
1.098	2-O-CH ₃	-	6-O-CH ₃		

- 16 -

Cmpd	R ₄	R ₄	R ₄	R ₅
1.099	2-O-CH ₃	4-CO-O-CH ₃	6-O-CH ₃	
1.100	2-CH ₃	4-Cl	6-CH ₃	
1.101	2-CH ₃	4-F	6-CH ₃	
1.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
1.103	2-F	4-Cl	5-O-i-C ₃ H ₇	
1.104	2-C1	4-Cl	5-O-CH ₃	
1.105		4-Cl	5-O-CH ₃	
1.106	2-F	4-Cl	5-CO-O-CH ₃	
1.107	2-F	4-Cl	5-CO-O-C ₂ H ₅	
1.108		4-Cl	5-CO-O-CH ₃	
1.109	2-Cl	4-Cl	5-CO-O-i-C ₃ H ₇	
1.110				4 - 0 - 🕒
1.111				4 - 0 — CI
1.112				4 - 0 — F
1.113				4-0-CF ₃
1.114	2-CH ₃			4-0-
1.115	;			4 - S —
1.110	5			4 - S — CI
1.117	7			4 - CH ₂ —
1.11	8			4 -CH ₂ -CI
1.11	9			4-CH ₂ -F

- 17 -

Cmpd	R ₄	R ₄	R ₄	R ₅
1.120				4-CH ₂ -CF ₃
1.121				4 - N-CHO
1.122	2-F	4-Cl	5-O-CH ₂ -CH=CH ₂	
1.123	2-F	4-Cl	5-O-CH ₂ -C≡CH	
1.124	2-Br			
1.125	2-CF ₃			
1.126	2-OCH ₃			
1.127	2-CH ₃			4.0
1.128	2-CH ₃			4-0-CF ₃
1.129	2-CH ₃			4 -0 -CI
1.130	2-CH ₃		6-CH ₃	4-0-
1.131	2-CH ₃		6-CH ₃	4-0-CF ₃
1.132	2-CH ₃		6-CH ₃	4-0-CI
1.133	2-CH ₃		6-CH ₃	4 -o — C1
1.134	2-CH₃	4-Br	6-CH ₃	Ġ
1.135		6-C ₂ H ₅	-	
1.136	2-C ₂ H ₅			
1.137	2-CH ₃	4-OC ₂ H ₅	6-CH ₃	
1.138	-	4-O-i-C ₃ H ₇	6-CH ₃	
1.139	2-CH ₃	4-O-n-C ₃ H ₇	6-CH ₃	
1.140	2-CH ₃	4-O-n-C ₁₀ H ₂₁	6-CH ₃	
1.141	2-CH ₃	4-O-n-C ₃ H ₇		

- 18 -

Cmpd	R ₄	R ₄	R ₄	R ₅
1.143	•	4-O-(CH ₂) ₂ OCH ₃ 4-O-(CH ₂) ₂ OCH ₃	6-CH ₃	
1.144 1.145	_	4-O-(CH ₂) ₂ OCH ₃ 4-O-n-C ₆ H ₁₃	6-CH₃	
1.146	2-CH ₃	V	·	4-0-
1.147	2-CH ₃		6-CH ₃	4-0-
1.148	2-CH ₃			4-0-CF ₃
1.149	2-CH ₃		6-CH ₃	4 -0 -CF ₃
1.150	2-CH ₃			4 - 0 - CF ₃
1.151	2-CH ₃		6-CH ₃	$4 - 0 - \bigvee_{C_1}^{N} - CF_3$
1.152	2-CH ₃			$5 \cdot 0 - \bigvee_{C1}^{N} - CF_3$
1.153	2-CH ₃			5 -0 — CF ₃
1.154	2-CH ₃		6-CH ₃	4 -s —
1.155	2-CH ₃		6-CH₃	4 - s — Ci
1.156	2-C ₂ H ₅		6-CH ₃	4 - s — \\\\
1.157 1.158 1.159	2-C ₂ H ₅ 2-C ₂ H ₅ 2-CH(CH ₃)	4-C ₂ H ₅	6-C ₂ H ₅	

- 19 -

Table 2: Compounds of formula Ie

$$\begin{array}{c|c}
 & O \\
 & N \\
 & N \\
 & N \\
\end{array}$$
(Ie)

Cmpd	R_4	R ₄	
2.001	3-Cl		
2.002	3-F		•
2.003	3-CH ₃		
2.004	5-C1		+
2.005	5-CF ₃		
2.006	3-Cl	5-Cl	
2.007	3-Cl	5-F	
2.008	3-Cl	5-CF ₃	
2.009	3-C1	5-NO ₂	
2.010	3-C1	5-SO ₂ -CH ₃	
2.011	3-F	5-F	
2.012	3-F	5-C1	
2.013	3-F	5-CF ₃	
2.014	3-NO ₂	5-NO ₂	
2.015	3-NO ₂	5-C1	
2.016	3-NO ₂	5-CF ₃	
2.017	3-CF ₃	5-Cl	
2.018	3-CF ₃	5-CF ₃	
2.019	3-CH ₃	5-CH ₃	

- 20 -

Table 3: Compounds of formula If

$$\begin{array}{c|c}
 & O \\
 & N \\
 & N \\
 & O
\end{array}$$

$$\begin{array}{c}
 & (R_4)_n \\
 & (R_5)_m
\end{array}$$
(If)

		O			
Cmpd	R ₄	R ₄	R ₄	R ₅	
3.001		Н			
3.002		2-CH ₃			
3.003		4-CH ₃			
3.004		2-CH ₃	4-CH ₃		
3.005		$2-CH_3$	6-CH ₃		
3.006		$2-CH_3$	5-CH ₃		
3.007		3-CH ₃	5-CH ₃		
3.008		2-CH ₃	3-CH ₃		
3.009		3-CH ₃	4-CH ₃		
3.010	2-CH ₃	4-CH ₃	6-CH ₃		
3.011	2-CH ₃	4-CH ₃	5-CH ₃		
3.012		2-C1			
3.013		4-C1			
3.014		2-C1	4-C1		
3.015		2-Cl	6-C1		
3.016		2-C1	6-F		
3.017		2-CH ₃	4-C1		
3.018		2-CH ₃	4-F		
3.019		2-C1	4-CH ₃		
3.020		2-C1	6-CH ₃		
3.021		2-F	4-F		
3.022		2-F	6-F		
3.023		2-CH ₃	4-O-CH ₃		
3.024		2-CH ₃	6-O-CH ₃		
3.025		2-C1	4-O-CH ₃		
3.026		2-Cl	6-O-CH ₃		

- 21 -

Cmpd R ₄	R ₄	R ₄	R ₅	
3.027	3-OCH ₃	4-OCH₃		
3.028	2-OCH ₃	5-OCH ₃		
3.029	2-OCH ₃	4-OCH ₃		
3.030	2-OCH ₃	6-OCH ₃		
3.031	2-CF ₃	6-CF ₃		
3.032	2-CF ₃	4-CF ₃		
3.033	3-CF ₃	5-CF ₃		
3.034	2-C1	4-CF ₃		
3.035	2-C1	6-CF ₃		
3.036	2-NO ₂	4-NO ₂		
3.037	2-Cl	4-NO ₂		
3.038	2-CH ₃	4-NO ₂		
3.039	2-O-CH ₃	4-NO ₂		
3.040	2-F	6-NO ₂		
3.041	2-Cl	6-NO ₂		
3.042	2-CH ₃	6-NO ₂		
3.043	2-O-CH ₃	6-NO ₂		
3.044	2-F	4-NO ₂		
3.045	2-CH ₃	$4-N(C_2H_5)_2$		
3.046	2-Cl	4-SO ₂ -CH ₃		
3.047	2-C1	4-SO-CH ₃		
3.048	2-C1	4-S-CH ₃		
3.049	2-Cl	6-SO ₂ -CH ₃		
3.050	2-Cl	6-SO-CH ₃		
3.051	2-C1	6-S-CH ₃		
3.052	2-CH ₃	4-SO ₂ -CH ₃		
3.053	2-CH ₃	4-SO-CH ₃		
3.054	2-CH ₃	4-S-CH ₃		
3.055	2-CH ₃	6-SO ₂ -CH ₃		
3.056	2-CH ₃	6-SO-CH ₃		
3.057	2-CH ₃	6-S-CH ₃		
3.058	2-O-CH ₃	6-SO ₂ -CH ₃		
3.059	2-O-CH ₃	6-SO-CH ₃		
3.060	2-O-CH ₃	6-S-CH ₃		
3.061	2-O-CH ₃	4-SO ₂ -CH ₃		
3.062	2-O-CH ₃	4-SO-CH ₃		
	-	•		

- 22 -

Cmpd	R ₄	R ₄	R ₄	R ₅	
3.063		2-O-CH ₃	4-S-CH ₃		
3.064		2-CH ₃	$6-N(C_2H_5)_2$		
3.065		2-Cl	$6-N(CH_3)_2$		
3.066		2-C1	4-N(CH ₃) ₂		
3.067		2-Cl	4-CO ₂ CH ₃		
3.068		2-CH ₃	$6-CO_2C_2H_5$		
3.069		2-CH ₃	$4-CO_2C_2H_5$		
3.070		2-CH ₃	4-CN		
3.071		2-CH ₃	6-CN		
3.072		2-C1	4-CN		
3.073		2-C1	6-CN		
3.074		2-Cl	4-CO-CH ₃		
3.075		2-O-CHF ₂	4-O-CHF ₂		
3.076		2-CH ₃	4-O-CHF ₂		
3.077		2-C1	4-O-CF ₃		
3.078		2-O-CF ₃	4-O-CH ₃		
3.079		2-O-CHF ₂	4-Cl		
3.080		2-O-CHF ₂	6-CH ₃		
3.081		2-O-CHF ₂	6-C1		
3.082	2-O-CHF ₂	4-CH ₃	6-CH ₃		
3.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃		
3.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇		
3.085	2-CH ₃	4-O-CH ₃	6-CH₃		
3.086	_	4-CF ₃	6-Cl		
3.087		4-CF ₃	6-F		
3.088		4-NO ₂	6-C1		
3.089		4-CI	6-C1		
3.090		4-F	6-F		
3.091		4-NO ₂	6-CH ₃		
3.092	_	4-C1	6-CH ₃		
3.093		4-O-CH ₃	6-C1		
3.094		4-C1	6-O-CH ₃		
3.095		4-O-CH ₃	6-F		
3.096		4-CH ₃	6-O-CH ₃		
3.097	_		6-CH ₃		
3.09	_		6-O-CH ₃		

- 23 -

Cmpd	R ₄	R ₄	R ₄	R ₅
3.099	2-O-CH ₃	4-CO-O-CH ₃	6-O-CH₃	
3.100	2-CH ₃	4-C1	6-CH ₃	
3.101	2-CH ₃	4-F	6-CH ₃	
3.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
3.103	2-F	4-C1	5-O-i-C ₃ H ₇	
3.104	2-C1	4-C1	5-O-CH ₃	
3.105		4-C1	5-O-CH ₃	
3.106	2-F	4-C1	5-CO-O-CH ₃	
3.107	2-F	4-C1	5-CO-O-C ₂ H ₅	
3.108		4-C1	5-CO-O-CH ₃	
3.109	2-C1	4-Cl	5-CO-O-i-C ₃ H ₇	
3.110				4 - 0 -
3.111				4 - 0 — CI
3.112				4 - 0 — F
3.113				4-0-CF ₃
3.114	2-CH ₃			4-0-
3.115				4 - 8 -
3.116				4 - s — C1
3.117				4 - CH ₂ —
3.118				4- CH ₂ -CI
3.119				4 CH ₂ — F

Cmpd	R ₄	R ₄	R ₄	R ₅
3.120				4-CH ₂ -CF ₃
3.121				4 - N — CHO
3.122	2-F	4-Cl	5-O-CH ₂ -CH=CH ₂	
3.123	2-F	4-Cl	5-O-CH ₂ -C≡CH	
3.124	2-Br			
3.125	2-CF ₃			
3.126				
	•			CI
3.127	2-CH ₃			4 -0 -CI
3.128	2-CH ₃			4-0-CF ₃
3.129	2-CH ₃			4 -0 -C1
3.130	2-CH ₃		6-CH ₃	4-0-
3.131	2-CH ₃		6-CH ₃	4-0-CF ₃
3.132	2-CH ₃		6-CH ₃	4-0-CI
3.133	2-CH ₃		6-CH₃	4 -0 — CI
3.134	2-CH ₃	4-Br	6-CH₃	ča
3.135	2-CH ₃	6-C ₂ H ₅		
3.136				
3.137	-		6-CH ₃	
3.138	-	4-O-i-C ₃ H ₇	6-CH ₃	
3.139	_	4-O-n-C ₃ H ₇	6-CH ₃	
3.140	2-CH ₃	4-O-n-C ₁₀ H ₂₁	6-CH ₃	

Cmpd	R ₄	R ₄	R ₄	R ₅
3.141	2-CH ₃	4-O-n-C ₃ H ₇		
3.142	2-CH ₃	4-O-(CH ₂) ₂ OCH ₃		
3.143	-	4-O-(CH ₂) ₂ OCH ₃	6-CH ₃	
3.144	2-CH ₃	4-O-(CH ₂) ₂ OCH ₃		
3.145	2-CH ₃	4-O-n-C ₆ H ₁₃	6-CH ₃	N.
3.146	2-CH ₃			4-0-
3.147	2-CH ₃		6-CH ₃	4-0-N-
3.148	2-CH ₃		·	4-0-N-CF ₃
3.149	2-CH ₃		6-CH ₃	4 -0 - CF ₃
3.150	2-CH ₃			4-0-CF ₃
3.151	2-CH ₃		6-CH ₃	Cí 4-0
3.152	2-CH ₃			5-0-K-CF ₃
3.153	2-CH ₃			5-0-(N-)-CF ₃
3.154	2-CH ₃		6-CH ₃	4 -s —
3.155	2-CH ₃		6-CH ₃	4 - s — C1
3.156	2-C ₂ H ₅		6-CH ₃	4 - s —

Table 4: Compounds of formula Ig

$$\begin{array}{c|c}
 & O \\
 & N \\$$

Cmpd	R ₄	R ₄	R ₄	R ₅	
4.001		Н			
4.002		2-CH ₃			
4.003		4-CH ₃			
4.004		2-CH ₃	4-CH ₃		
4.005		2-CH ₃	6-CH ₃		
4.006		2-CH ₃	5-CH ₃		
4.007		3-CH ₃	5-CH ₃	·	
4.008		2-CH ₃	3-CH ₃		
4.009		3-CH ₃	4-CH ₃		
4.010	2-CH ₃	4-CH ₃	6-CH ₃		
4.011	2-CH ₃	4-CH ₃	5-CH ₃		
4.012		2-Cl			
4.013		4-C1			
4.014		2-C1	4-C1		
4.015		2-C1	6-C1		
4.016		2-C1	6-F	•	
4.017		2-CH ₃	4-C1		
4.018		2-CH ₃	4-F		
4.019		2-C1	4-CH ₃		
4.020		2-C1	6-CH ₃		
4.021		2-F	4-F		
4.022		2-F	6-F		
4.023		2-CH ₃	4-O-CH ₃		
4.024		2-CH ₃	6-O-CH ₃		
4.025		2-C1	4-O-CH ₃		
4.026		2-C1	6-O-CH ₃		
4.027		3-OCH ₃	4-OCH ₃		

- 27 -

Cmpd R ₄	R ₄	R ₄	R ₅	
4.028	2-OCH ₃	5-OCH ₃		
4.029	2-OCH ₃	4-OCH ₃		
4.030	2-OCH ₃	6-OCH₃		
4.031	2-CF ₃	6-CF ₃		
4.032	2-CF ₃	4-CF ₃		
4.033	3-CF ₃	5-CF ₃		
4.034	2-C1	4-CF ₃		
4.035	2-C1	6-CF ₃		
4.036	2-NO ₂	4-NO ₂	,	
4.037	2-C1	4-NO ₂		
4.038	2-CH ₃	4-NO ₂		
4.039	2-O-CH ₃	4-NO ₂		
4.040	2-F	6-NO ₂		
4.041	2-C1	6-NO ₂		
4.042	2-CH ₃	6-NO ₂		
4.043	2-O-CH ₃	6-NO ₂		
4.044	2-F	4-NO ₂		
4.045	2-CH ₃	$4-N(C_2H_5)_2$		
4.046	2-C1	4-SO ₂ -CH ₃		
4.047	2-C1	4-SO-CH ₃		
4.048	2-Cl	4-S-CH ₃		
4.049	2-C1	6-SO ₂ -CH ₃		
4.050	2-C1	6-SO-CH ₃		
4.051	2-Cl	6-S-CH ₃		
4.052	2-CH ₃	4-SO ₂ -CH ₃		
4.053	2-CH ₃	4-SO-CH ₃		
4.054	2-CH ₃	4-S-CH ₃		
4.055	2-CH ₃	6-SO ₂ -CH ₃		
4.056	2-CH ₃	6-SO-CH ₃		
4.057	2-CH ₃	6-S-CH ₃		
4.058	2-O-CH ₃	6-SO ₂ -CH ₃		
4.059	2-O-CH ₃	6-SO-CH ₃		
4.060	2-O-CH ₃	6-S-CH ₃		
4.061	2-O-CH ₃	4-SO ₂ -CH ₃		
4.062	2-O-CH ₃	4-SO-CH ₃		
4.063	2-O-CH ₃	4-S-CH ₃		

- 28 -

4.

Cmpd	R ₄	R ₄	R ₄	R ₅
4.064		2-CH ₃	6-N(C ₂ H ₅) ₂	
4.065		2-C1	$6-N(CH_3)_2$	
4.066		2-Cl	$4-N(CH_3)_2$	
4.067		2-Cl	4-CO ₂ CH ₃	
4.068		2-CH ₃	$6-CO_2C_2H_5$	
4.069		2-CH ₃	$4-CO_2C_2H_5$	
4.070		2-CH ₃	4-CN	
4.071		2-CH ₃	6-CN	
4.072		2-Cl	4-CN	
4.073		2-C1	6-CN	
4.074		2-Cl	4-CO-CH ₃	
4.075		2-O-CHF ₂	4-O-CHF ₂	
4.076		2-CH ₃	4-O-CHF ₂	
4.077		2-Cl	4-O-CF ₃	
4.078		2-O-CF ₃	4-O-CH ₃	
4.079		2-O-CHF ₂	4-C1	
4.080		2-O-CHF ₂	6-CH ₃	
4.081		2-O-CHF ₂	6-C1	
4.082	2-O-CHF ₂	4-CH ₃	6-CH ₃	
4.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃	
4.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇	
4.085	2-CH ₃	4-O-CH ₃	6-CH ₃	
4.086	2-Cl	4-CF ₃	6-C1	
4.087	2-C1	4-CF ₃	6-F	
4.088	2-C1	4-NO ₂	6-C1	
4.089	2-C1	4-Cl	6-C1	
4.090	2-F	4-F	6-F	
4.091	2-CH ₃	4-NO ₂	6-CH ₃	
4.092	2-C1	4-Cl	6-CH ₃	
4.093	2-C1	4-O-CH ₃	6-C1	
4.094	2-C1	4-C1	6-O-CH ₃	
4.095	2-F	4-O-CH ₃	6-F	
4.096	2-O-CH ₃	4-CH ₃	6-O-CH ₃	
4.097	2-O-CH ₃		6-CH ₃	
4.098	_		6-O-CH ₃	
4.099	•		6-O-CH ₃	
	_			

- 29 -

Cmpd	R ₄	R ₄	R ₄ .	R ₅
4.100	2-CH ₃	4-Cl	6-CH ₃	
4.101	2-CH ₃	4-F	6-CH ₃	
4.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
4.103	2-F	4-C1	5-O-i-C ₃ H ₇	
4.104	2-Cl	4-C1	5-O-CH ₃	
4.105		4-Cl	5-O-CH ₃	
4:106	2-F	4-C1	5-CO-O-CH ₃	
4.107	2-F	4-C1	5-CO-O-C ₂ H ₅	
4.108		4-Cl	5-CO-O-CH ₃	
4.109	2-C1	4-Cl	5-CO-O-i-C ₃ H ₇	
4.110				4 - 0
4.111				4 - 0 — CI
4.112				4 - 0 — F
4.113				4-0-CF3
4.114	2-CH ₃			4-0-
4.115				4 - S —
4.116				4 - S — C1
4.117				4 - CH ₂
4.118				4-CH ₂ -CI
4.119				4-CH ₂ -F
4.120				4-CH ₂ -CF ₃

WO 96/11574

2199636

PCT/EP95/03935

Cmpd	R ₄	R ₄	R ₄	R ₅	
4.121				4 - N-(CHO)	
4.122	2-F	4-C1	5-O-CH ₂ -CH=CH ₂		
4.123	2-F	4-Cl	5-O-CH ₂ -C=CH		

- 31 -

Table 5: Compounds of formula Ih

$$\begin{array}{c|c}
 & O \\
 & N \\
 & N \\
 & O \\$$

Cpmd	R ₄	R ₄	R ₄	R ₅	
5.001		Н			
5.002		2-CH ₃			
5.003		4-CH ₃			
5.004		2-CH ₃	4-CH ₃		
5.005		2-CH ₃	6-CH ₃		
5.006		2-CH ₃	5-CH ₃		
5.007		3-CH ₃	5-CH ₃		
5.008		2-CH ₃	3-CH ₃		<u>.</u>
5.009		3-CH ₃	4-CH ₃		·
5.010	2-CH ₃	4-CH ₃	6-CH ₃		
5.011	2-CH ₃	4-CH ₃	5-CH ₃		
5.012		2-Cl		,	
5.013		4-C1			
5.014		2-C1	4-CI		
5.015		2-Cl	6-CI		
5.016		2-C1	6-F		
5.017		2-CH ₃	4-C1		
5.018		2-CH ₃	4-F		
5.019		2-CI	4-CH ₃		
5.020		2-C1	6-CH ₃		
5.021		2-F	4-F		
5.022		2-F	6-F		
5.023		2-CH ₃	4-O-CH ₃		
5.024		2-CH ₃	6-O-CH ₃		
5.025		2-C1	4-O-CH ₃		
5.026		2-C1	6-O-CH ₃		

PCT/EP95/03935

-32-

-32-				
Cmpd R ₄	R ₄	R ₄	R ₅	
5.027	3-OCH ₃	4-OCH₃		
5.028	2-OCH ₃	5-OCH ₃		
5.029	2-OCH ₃	4-OCH ₃		
5.030	2-OCH ₃	6-OCH ₃		
5.031	2-CF ₃	6-CF ₃		
5.032	2-CF ₃	4-CF ₃		
5.033	3-CF ₃	5-CF ₃		
5.034	2-Cl	4-CF ₃		
5.035	2-C1	6-CF ₃		
5.036	2-NO ₂	4-NO ₂		
5.037	2-Cl	4-NO ₂		
5.038	2-CH ₃	4-NO ₂		
5.039	2-O-CH ₃	4-NO ₂		
5.040	2-F	6-NO ₂		
5.041	2-C1	6-NO ₂		
5.042	2-CH ₃	6-NO ₂		
5.043	2-O-CH ₃	6-NO ₂		
5.044	2-F	4-NO ₂		
5.045	2-CH ₃	$4-N(C_2H_5)_2$		
5.046	2-Cl	4-SO ₂ -CH ₃		
5.047	2-C1	4-SO-CH ₃		
5.048	2-C1	4-S-CH ₃		
5.049	2-Cl	6-SO ₂ -CH ₃		
5.050	2-C1	6-SO-CH ₃		
5.051	2-C1	6-S-CH ₃		
5.052	2-CH ₃	4-SO ₂ -CH ₃		
5.053	2-CH ₃	4-SO-CH ₃		
5.054	2-CH ₃	4-S-CH ₃		
5.055	2-CH ₃	6-SO ₂ -CH ₃		
5.056	2-CH ₃	6-SO-CH ₃		
5.057	2-CH ₃	6-S-CH ₃		
5.058	2-O-CH₃	6-SO ₂ -CH ₃		
5.059	2-O-CH ₃	6-SO-CH ₃		
5.060	2-O-CH ₃	6-S-CH ₃		
5.061	2-O-CH ₃	4-SO ₂ -CH ₃		
5.062	2-O-CH ₃	4-SO-CH ₃		

-33-

Cmpd	R ₄	R ₄	R_4	R ₅	
5.063		2-O-CH ₃	4-S-CH ₃		
5.064		2-CH ₃	$6-N(C_2H_5)_2$		
5.065		2-C1	6-N(CH ₃) ₂		
5.066		2-C1	4-N(CH ₃) ₂		
5.067		2-C1	4-CO ₂ CH ₃		
5.068		2-CH ₃	6-CO ₂ C ₂ H ₅		
5.069		2-CH ₃	4-CO ₂ C ₂ H ₅		
5.070		2-CH ₃	4-CN		
5.071		2-CH ₃	6-CN		
5.072		2-C1	4-CN		
5.073		2-C1	6-CN		
5.074		2-C1	4-CO-CH ₃		
5.075		2-O-CHF ₂	4-O-CHF ₂		
5.076		2-CH ₃	4-O-CHF ₂		
5.077		2-Cl	4-O-CF ₃		
5.078		2-O-CF ₃	4-O-CH ₃	•	
5.079		2-O-CHF ₂	4-C1		•
5.080		2-O-CHF ₂	6-CH ₃		
5.081		2-O-CHF ₂	6-C1		
5.082	2-O-CHF ₂	4-CH ₃	6-CH ₃		
5.083	2-CH ₃	$4-t-C_4H_9$	6-CH ₃		
5.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇		
5.085	2-CH ₃	4-O-CH ₃	6-CH ₃		
5.086	2-Cl	4-CF ₃	6-C1		
5.087	2-C1	4-CF ₃	6-F		
5.088	2-CI	4-NO ₂	6-C1		
5.089	2-C1	4-Cl	6-C1		
5.090	2-F	4-F	6-F		
5.091	2-CH ₃	4-NO ₂	6-CH ₃		
5.092	2-Cl	4-Cl	6-CH ₃		
5.093	2-Cl	4-O-CH ₃	6-Cl		
5.094	2-C1	4-C1	6-O-CH ₃		
5.095	2-F	4-O-CH ₃	6-F		
5.096	2-O-CH ₃	4-CH ₃	6-O-CH ₃		
5.097	2-O-CH ₃	4-O-CH ₃	6-CH ₃		
5.098	2-O-CH ₃	4-O-CH ₃	6-O-CH ₃		

_	7/
-	JЧ

Cmpd	R ₄	R ₄	R ₄	R₅
5.099	2-O-CH ₃	4 CO O CT		
5.100	2-CH ₃	4-CO-O-CH₃ 4-Cl	6-O-CH ₃	
5.101	2-CH ₃	4-F	6-CH ₃	
5.102	2-CH ₃	4-CH ₃	6-CH ₃	
5.103	2-F	4-Cl	6-O-CH ₃	
5.104	2-C1	4-C1	5-O-i-C ₃ H ₇	
5.105	- 0.	4-C1	5-O-CH ₃	
5.106	2-F	4-Cl	5-O-CH ₃	
5.107	2-F	4-Cl	5-CO-O-CH ₃	
5.108		4-C1	5-CO-O-C ₂ H ₅	
5.109	2-C1	4-CI	5-CO-O-CH ₃	
-1		4 C1	5-CO-O-i-C ₃ H ₇	_
5.110				4-0-
5.111				4 - 0 — CI
5.112				4 - 0 - F
5.113				4-0-CF ₃
5.114	2-CH ₃			4-0-
5.115				4 - S —
5.116				4 - s — CI
5.117				4 - CH ₂
5.118				4 -CH ₂ -CI
5.119				4 -CH ₂

WO 96/11574

PCT/EP95/03935

- 35 -

Cmpd	R ₄	R ₄	R ₄	R ₅
5.120				4 CH ₂ - CF ₃
5.121				4 - N - CHO
5.122	2-F	4-Cl	5-O-CH ₂ -CH=CH ₂	
5.123	2-F	4-C1	5-O-CH ₂ -C≡CH	

- 36 -

Table 6: Compounds of formula Ii

(Ii)

Cmpd	R ₄	R ₄	R ₄	R ₅	
6.001		Н			
6.002		2-CH ₃			
6.003		4-CH ₃			
6.004		2-CH ₃	4-CH ₃		
6.005		2-CH ₃	6-CH ₃		
6.006		2-CH ₃	5-CH ₃		
6.007		3-CH ₃	5-CH ₃		
6.008		2-CH ₃	3-CH ₃		
6.009		3-CH ₃	4-CH ₃		
6.010	2-CH ₃	4-CH ₃	6-CH ₃		
6.011	2-CH ₃	4-CH ₃	5-CH ₃		
6.012		2-Cl			
6.013		4-C1			
6.014		2-C1	4-C1		
6.015		2-Cl	6-C1		
6.016		2-Cl	6-F		
6.017		2-CH ₃	4-C1		
6.018		2-CH ₃	4-F		
6.019		2-Cl	4-CH ₃	·	
6.020		2-CI	6-CH ₃		
6.021		2-F	4-F		
6.022		2-F	6-F		
6.023		$2-CH_3$	4-O-CH ₃		
6.024		2-CH ₃	6-O-CH ₃		
6.025		2-Cl	4-O-CH ₃		
6.026		2-Cl	6-O-CH ₃		
6.027		3-OCH ₃	4-OCH ₃		

- 37 -

Cmpd R ₄	R ₄	R ₄	R ₅	
6.028	2-OCH ₃	5-OCH ₃		
6.029	2-OCH ₃	4-OCH ₃		
6.030	2-OCH ₃	6-OCH₃		
6.031	2-CF ₃	6-CF ₃		
6.032	2-CF ₃	4-CF ₃		
6.033	3-CF ₃	5-CF ₃		
6.034	2-Cl	4-CF ₃		
6.035	2-Cl	6-CF ₃		
6.036	2-NO ₂	4-NO ₂		
6.037	2-C1	4-NO ₂		
6.038	2-CH ₃	4-NO ₂		
6.039	2-O-CH ₃	4-NO ₂		
6.040	2-F	6-NO ₂		
6.041	2-C1	6-NO ₂		
6.042	2-CH ₃	6-NO ₂		•
6.043	2-O-CH ₃	6-NO ₂		
6.044	2-F	4-NO ₂		
6.045	2-CH ₃	$4-N(C_2H_5)_2$		
6.046	2-C1	4-SO ₂ -CH ₃		
6.047	2-Cl	4-SO-CH ₃		
6.048	2-C1	4-S-CH ₃		
6.049	2-C1	6-SO ₂ -CH ₃		
6.050	2-C1	6-SO-CH ₃		
6.051	2-C1	6-S-CH₃		
6.052	2-CH ₃	4-SO ₂ -CH ₃		
6.053	2-CH ₃	4-SO-CH ₃		
6.054	2-CH ₃	4-S-CH ₃		
6.055	2-CH ₃	6-SO ₂ -CH ₃		
6.056	2-CH ₃	6-SO-CH ₃		
6.057	2-CH ₃	6-S-CH ₃		
6.058	2-O-CH ₃	6-SO ₂ -CH ₃		
6.059	2-O-CH ₃	6-SO-CH ₃		
6.060	2-O-CH ₃	6-S-CH ₃		
6.061	2-O-CH ₃	4-SO ₂ -CH ₃		
6.062	2-O-CH ₃	4-SO-CH ₃		

- 38 -

Cmpd	R ₄	R ₄	R ₄	R ₅	
6.063	7-7-	2-O-CH ₃	4-S-CH₃		
6.064		2-CH ₃	$6-N(C_2H_5)_2$		
6.065		2-Cl	6-N(CH ₃) ₂		
6.066		2-Cl	4-N(CH ₃) ₂		
6.067		2-Cl	4-CO ₂ CH ₃		
6.068		2-CH ₃	6-CO ₂ C ₂ H ₅		
6.069		2-CH ₃	4-CO ₂ C ₂ H ₅		
6.070		2-CH ₃	4-CN		
6.071		2-CH ₃	6-CN		
6.072		2-C1	4-CN		
6.073		2-C1	6-CN		
6.074		2-Cl	4-CO-CH ₃		
6.075		2-O-CHF ₂	4-O-CHF ₂		
6.076		2-CH ₃	4-O-CHF ₂		
6.077		2-C1	4-O-CF ₃		
6.078		2-O-CF ₃	4-O-CH ₃		
6.079		2-O-CHF ₂	4-C1		
6.080		2-O-CHF ₂	6-CH ₃		
6.081		2-O-CHF ₂	6-C1		
6.082	2-O-CHF ₂	4-CH ₃	6-CH ₃		
6.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃		
6.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇		
6.085	2-CH ₃	4-O-CH ₃	6-CH ₃		
6.086	2-C1	4-CF ₃	6-C1		
6.087	2-C1	4-CF ₃	6-F		<i>-</i> .
6.088	2-Cl	4-NO ₂	6-C1		
6.089	2-Cl	4-Cl	6-C1		
6.090	2-F	4-F	6-F		
6.091	2-CH ₃	4-NO ₂	6-CH ₃		
6.092	2-C1	4-C1	6-CH ₃		
6.093	2-C1	4-O-CH ₃	6-C1		
6.094	2-Cl	4-CI	6-O-CH₃		
6.095	2-F	4-O-CH ₃	6-F		
6.096	2-O-CH ₃	4-CH ₃	6-O-CH₃		
6.097	2-O-CH ₃	4-O-CH ₃	6-CH ₃		
6.098	2-O-CH ₃	4-O-CH ₃	6-O-CH ₃		

- 39 -

Cmpd	R ₄	R ₄	R ₄	R ₅
-				
6.099	2-O-CH ₃	4-CO-O-CH ₃	6-O-CH ₃	
6.100	2-CH ₃	4-C1	6-CH ₃	
6.101	2-CH ₃	4-F	6-CH ₃	
6.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
6.103	2-F	4-C1	5-O-i-C ₃ H ₇	
6.104	2-Cl	4-Cl	5-O-CH ₃	
6.105		4-Cl	5-O-CH ₃	
6.106	2-F	4-Cl	5-CO-O-CH ₃	
6.107	2-F	4-Cl	5-CO-O-C ₂ H ₅	
6.108		4-Cl	5-CO-O-CH ₃	
6.109	2-Cl	4-Cl	5-CO-O-i-C ₃ H ₇	
6.110				4 - 0 -
6.111				4 - 0 — C1
6.112				4 - 0 - F
6.113				4-0-CF3
6.114	2-CH ₃			4-0-
6.115				4 - 8 -
6.116				4 - S — CI
6.117				4 - CH ₂
6.118				4 -CH ₂ -CI
6.119				4 -CH ₂ - F

- 40 -

Cmpd	R ₄	R ₄	R ₄	R ₅
6.120				4.CH ₂ -CF ₃
6.121				4 - N - CHO
6.122	2-F	4-Cl	5-O-CH ₂ -CH=CH ₂	
6.123	2-F	4-Cl '	5-O-CH ₂ -C≡CH	
6.124	2-Br			
6.125	2-CF ₃			
6.126	2-OCH ₃			
				CI \
6.127	2-CH ₃			4 -o - C1
6.128	2-CH ₃			4-0-CF3
6.129	2-CH ₃			4-0-CI
6.130	2-CH ₃		6-CH ₃	4-0-
6.131	2-CH ₃		6-CH ₃	4-0-CF ₃
6.132	2-CH ₃		6-CH ₃	4 -0 -C1
6.133	2-CH ₃		6-CH ₃	4 -0 -C1
	o 677	4 D-	6 CU	ď
6.134	_	4-Br	6-CH₃	
6.135	_	6-C ₂ H ₅		
6.136		6-C ₂ H ₅ 4-OC ₂ H ₅	6-CH₃	
6.137	-	4-0-i-C ₃ H ₇	6-CH ₃	
6.138	=	4-O-n-C ₃ H ₇	6-CH ₃	
6.139	_	4-O-n-C ₁₀ H ₂₁	6-CH ₃	
6.140	2-CH ₃	4-0-11-0101121	ر کیدی	

- 41 -

Cmpd	R ₄	R ₄	R ₄	R ₅
6.141	2-CH ₃	4-O-n-C ₃ H ₇		
6.142	2-CH ₃	4-O-(CH ₂) ₂ OCH ₃		
6.143	-	4-O-(CH ₂) ₂ OCH ₃	6-CH ₃	
6.144	_	4-O-(CH ₂) ₂ OCH ₃		
6.145	2-CH ₃	4-O-n-C ₆ H ₁₃	6-CH ₃	N.
6.146	2-CH ₃			4-0-
6.147	2-CH ₃		6-CH ₃	4 -o - N
6.148	2-CH ₃			4 -0 - CF ₃
6.149	2-CH ₃		6-CH ₃	4 -0 - CF ₃
6.150	2-CH ₃			4-0-KP3
6.151	2-CH ₃		6-CH ₃	4-0-X
6.152	2-CH ₃			5-0-KP3
6.153	2-CH ₃			5-o-N
6.154	2-CH ₃		6-CH ₃	4 - S —
6.155	2-CH ₃		6-CH ₃	4 - s — CI
6.156	2-C ₂ H ₅		6-CH ₃	4 -s —

- 42 -

Table 7: Compounds of formula Ij

- 43 -

Cmpd R ₄	R ₄	R ₄	R ₅	
7.028	2-OCH ₃	5-OCH ₃		
7.029	2-OCH ₃	4-OCH ₃		
7.030	2-OCH ₃	6-OCH ₃		
7.031	2-CF ₃	6-CF ₃		
7.032	2-CF ₃	4-CF ₃		
7.033	3-CF ₃	5-CF ₃		
7.034	2-C1	4-CF ₃		
7.035	2-Cl	6-CF ₃		
7.036	2-NO ₂	4-NO ₂		
7.037	2-Cl	4-NO ₂		
7.038	2-CH ₃	4-NO ₂		
7.039	2-O-CH ₃	4-NO ₂		
7.040	2-F	6-NO ₂		
7.041	2-Ci	6-NO ₂		
7.042	2-CH ₃	6-NO ₂	•	
7.043	2-O-CH ₃	6-NO ₂		
7.044	2-F	4-NO ₂		
7.045	2-CH ₃	$4-N(C_2H_5)_2$		
7.046	2-C1	4-SO ₂ -CH ₃		
7.047	2-C1	4-SO-CH ₃		
7.048	2-C1	4-S-CH ₃		
7.049	2-C1	6-SO ₂ -CH ₃		
7.050	2-C1	6-SO-CH ₃		
7.051	2-C1	6-S-CH ₃		
7.052	2-CH ₃	4-SO ₂ -CH ₃		
7.053	2-CH ₃	4-SO-CH ₃		
7.054	2-CH ₃	4-S-CH ₃		
7.055	2-CH ₃	6-SO ₂ -CH ₃		
7.056	2-CH ₃	6-SO-CH ₃		
7.057	2-CH ₃	6-S-CH ₃		
7.058	2-O-CH ₃	6-SO ₂ -CH ₃		
7.059	2-O-CH ₃	6-SO-CH ₃		
7.060	2-O-CH ₃	6-S-CH ₃		
7.061	2-O-CH ₃	4-SO ₂ -CH ₃		
7.062	2-O-CH ₃	4-SO-CH ₃		
		. 50 043		

- 44 -

Cmpd	R ₄	R ₄	R ₄	R ₅	
7.063		2-O-CH ₃	4-S-CH ₃		
7.064		2-CH ₃	$6-N(C_2H_5)_2$		
7.065		2-Cl	6-N(CH ₃) ₂		
7.066		2-Cl	4-N(CH ₃) ₂		
7.067		2-Cl	4-CO ₂ CH ₃		
7.068		2-CH ₃	6-CO ₂ C ₂ H ₅		
7.069		2-CH ₃	4-CO ₂ C ₂ H ₅		
7.070		2-CH ₃	4-CN		
7.071		2-CH ₃	6-CN		
7.072		2-Cl	4-CN		
7.073		2-Cl	6-CN		
7.074		2-C1	4-CO-CH ₃		
7.075		2-O-CHF ₂	4-O-CHF ₂		
7.076		2-CH ₃	4-O-CHF ₂		
7.077		2-Cl	4-O-CF ₃		
7.078		2-O-CF ₃	4-O-CH ₃		
7.079		2-O-CHF ₂	4-Cl		
7.080		2-O-CHF ₂	6-CH ₃		
7.081		2-O-CHF ₂	6-C1		
7.082	2-O-CHF ₂	4-CH ₃	6-CH ₃		
7.083	2-CH ₃	$4-t-C_4H_9$	6-CH ₃		
7.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇		
7.085	2-CH ₃	4-O-CH ₃	6-CH ₃		
7.086	2-C1	4-CF ₃	6-C1		
7.087	2-C1	4-CF ₃	6-F		
7.088	2-Cl	4-NO ₂	6-C1		
7.089	2-C1	4-C1	6-C1		
7.090	2-F	4-F	6-F		
7.091	2-CH ₃	4-NO ₂	6-CH ₃		
7.092	2-Cl	4-C1	6-CH ₃		
7.093	2-C1	4-O-CH ₃	6-Cl		
7.094	2-C1	4-CI	6-O-CH ₃		
7.095	2-F	4-O-CH ₃	6-F		
7.096	2-O-CH ₃	4-CH ₃	6-O-CH ₃		
7.097	2-O-CH ₃	4-O-CH ₃	6-CH ₃		

Cmpd	R.	R ₄	R_4	R ₅
		**4	4	3
7.098	2-O-CH ₃	4-O-CH ₃	6-O-CH₃	
7.099	2-O-CH ₃	=	6-O-CH ₃	
7.100	2-CH ₃	4-C1	6-CH ₃	
7.101	2-CH ₃	4-F	6-CH ₃	
7.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
7.103	2-F	4-C1	5-O-i-C ₃ H ₇	
7.104	2-C1	4-C1	5-O-CH ₃	
7.105		4-C1	5-O-CH ₃	
7.106	2-F	4-C1	5-CO-O-CH ₃	
7.107	2-F	4-C1	5-CO-O-C ₂ H ₅	
7.108		4-C1	5-CO-O-CH ₃	
7.109	2-C1	4-C1	5-CO-O-i-C ₃ H ₇	
7.110				4 - 0
7.111				4 - 0 — CI
7.112				4 - O — F
7.113				4-0-CF3
7.114	2-CH ₃			4-0-
7.115				4 - S -
7.116				4 - s — Ci
7.117				4 - CH ₂ —
7.118				4 -CH ₂ -CI

2199636

PCT/EP95/03935

Cmpd	R ₄	R ₄	R ₄	R ₅
7.119				4 -CH ₂ - F
7.120				4 CH ₂ - CF ₃
7.121				4 - N — CHO
7.122	2-F	4-C1	5-O-CH ₂ -CH=CH ₂	
7.123	2-F	4-C1	5-O-CH ₂ -C≖CH	

- 47 -

Table 8: Compounds of formula Ik

$$\begin{array}{c|c}
 & O \\
 & N \\
 & N \\
 & O
\end{array}$$

$$\begin{array}{c}
 & (R_4)_n \\
 & (R_5)_m
\end{array}$$
(Ik)

Cmpd	R ₄	R ₄	R_4	R ₅	
8.001		Н			
8.002		2-CH ₃			
8.003		4-CH ₃			
8.004		2-CH ₃	4-CH ₃		
8.005		2-CH ₃	6-CH ₃		
8.006		2-CH ₃	5-CH ₃		
8.007		3-CH ₃	5-CH ₃		
8.008		2-CH ₃	3-CH ₃		
8.009		3-CH ₃	4-CH ₃		
8.010	2-CH ₃	4-CH ₃	6-CH ₃		
8.011	2-CH ₃	4-CH ₃	5-CH ₃		
8.012		2-C1			
8.013		4-C1			
8.014		2-C1	4-C1		
8.015		2-C1	6-C1		
8.016		2-Cl	6-F		
8.017		2-CH ₃	4-C1		
8.018		2-CH ₃	4-F		
8.019		2-Cl	4-CH ₃		
8.020		2-C1	6-CH ₃		
8.021		2-F	4-F		
8.022		2-F	6-F		
8.023		2-CH ₃	4-O-CH ₃		
8.024		2-CH ₃	6-O-CH ₃		
8.025		2-C1	4-O-CH ₃		
8.026		2-Cl	6-O-CH ₃		

- 48 -

Cmpd R ₄	R ₄	R ₄	R ₅	
8.027	3-OCH₃	4-OCH ₃		
8.028	2-OCH ₃	5-OCH ₃		
8.029	2-OCH ₃	4-OCH ₃		
8.030	2-OCH ₃	6-OCH ₃		
8.031	2-CF ₃	6-CF ₃		
8.032	2-CF ₃	4-CF ₃		
8.033	3-CF ₃	5-CF ₃		
8.034	2-C1	4-CF ₃		
8.035	2-C1	6-CF ₃		
8.036	2-NO ₂	4-NO ₂		
8.037	2-C1	4-NO ₂		
8.038	2-CH ₃	4-NO ₂		
8.039	2-O-CH ₃	4-NO ₂		
8.040	2-F	6-NO ₂		
8.041	2-C1	6-NO ₂		
8.042	2-CH ₃	6-NO ₂		
8.043	2-O-CH ₃	6-NO ₂		
8.044	2-F	4-NO ₂		
8.045	2-CH ₃	$4-N(C_2H_5)_2$		
8.046	2-C1	4-SO ₂ -CH ₃		
8.047	2-C1	4-SO-CH ₃		
8.048	2-C1	4-S-CH ₃		
8.049	2-C1	6-SO ₂ -CH ₃		
8.050	2-Cl	6-SO-CH ₃		
8.051	2-Cl	6-S-CH ₃		,
8.052	2-CH ₃	4-SO ₂ -CH ₃		
8.053	2-CH ₃	4-SO-CH ₃		
8.054	2-CH ₃	4-S-CH ₃		
8.055	2-CH ₃	6-SO ₂ -CH ₃		
8.056	2-CH ₃	6-SO-CH ₃		
8.057	2-CH ₃	6-S-CH ₃		
8.058	2-O-CH ₃	6-SO ₂ -CH ₃		
8.059	2-O-CH ₃	6-SO-CH ₃		
8.060	2-O-CH ₃	6-S-CH ₃		
8.061	2-O-CH ₃	4-SO ₂ -CH ₃		
8.062	2-O-CH ₃	4-SO-CH ₃		
-	3	•		

- 49 -

Cmpd	R ₄	R ₄	R ₄	R ₅
8.063		2-O-CH ₃	4-S-CH ₃	
8.064		2-CH ₃	$6-N(C_2H_5)_2$	
8.065		2-C1	6-N(CH ₃) ₂	
8.066		2-Cl	$4-N(CH_3)_2$	
8.067		2-C1	4-CO ₂ CH ₃	
8.068		2-CH ₃	6-CO ₂ C ₂ H ₅	
8.069		2-CH ₃	4-CO ₂ C ₂ H ₅	
8.070		2-CH ₃	4-CN	
8.071		2-CH ₃	6-CN	
8.072		2-Cl	4-CN	
8.073		2-Cl	6-CN	
8.074		2-Cl	4-CO-CH ₃	
8.075		2-O-CHF ₂	4-O-CHF ₂	
8.076		2-CH ₃	4-O-CHF ₂	
8.077		2-Cl	4-O-CF ₃	
8.078		2-O-CF ₃	4-O-CH ₃	
8.079		2-O-CHF ₂	4-Cl	
8.080		2-O-CHF ₂	6-CH ₃	
8.081		2-O-CHF ₂	6-Cl	
8.082	2-O-CHF ₂	4-CH ₃	6-CH ₃	
8.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃	
8.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇	
8.085	2-CH ₃	4-O-CH ₃	6-CH ₃	
8.086	2-C1	4-CF ₃	6-Cl	
8.087	2-C1	4-CF ₃	6-F	
8.088	2-C1	4-NO ₂	6-C1	
8.089	2-C1	4-C1	6-C1	
8.090	2-F	4-F	6-F	
8.091	2-CH ₃	4-NO ₂	6-CH ₃	
8.092	2-Cl	4-C1	6-CH ₃	
8.093	2-Cl	4-O-CH ₃	6-C1	
8.094	2-Cl	4-C1	6-O-CH ₃	
8.095	2-F	4-O-CH ₃	6-F	
8.096	2-O-CH ₃	_	6-O-CH ₃	
8.097	2-O-CH ₃	-	6-CH ₃	
8.098	•	4-O-CH ₃	6-O-CH ₃	

Cmpd	R ₄	R ₄	R ₄	R ₅
8.099	2-O-CH ₃	4-CO-O-CH ₃	6-O-CH ₃	
8.100	2-CH ₃	4-C1	6-CH ₃	
8.101	2-CH ₃	4-F	6-CH ₃	
8.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
8.103	2-F	4-C1	5-O-i-C ₃ H ₇	
8.104	2-C1	4-C1	5-O-CH ₃	
8.105		4-C1	5-O-CH ₃	
8.106	2-F	4-Cl	5-CO-O-CH ₃	
8.107	2-F	4-Cl	5-CO-O-C ₂ H ₅	
8.108		4-Cl	5-CO-O-CH ₃	
8.109	2-Cl	4-Cl	5-CO-O-i-C ₃ H ₇	
8.110				4 - 0
8.111				4 - 0 - 0
8.112				4 - O
8.113				4-0-CF3
8.114	2-CH ₃			4-0-
8.115				4 - S -
8.116				4 - s — CI
8.117				4 - CH ₂
8.118				4 -CH ₂
8.119				4.CH ₂ -F

- 51 -

Cmpd	R ₄	R ₄	R ₄	R ₅
8.120				4 - CH ₂ - CF ₃
8.121				4 - N-CHO
8.122	2-F	4-C1	5-O-CH ₂ -CH=CH ₂	
8.123	2-F	4-C1	5-O-CH ₂ -C≡CH	

- 52 -

Table 9: Compounds of formula II

$$\begin{array}{c|c}
 & O \\
 & N \\
 & N \\
 & O \\
 & (R_4)_n \\
 & (R_5)_m
\end{array}$$
(II)

Cmpd	R ₄	R ₄	R ₄	R ₅	
9.001		Н			
9.002		2-CH ₃			
9.003		4-CH ₃			
9.004		2-CH ₃	4-CH ₃		
9.005		2-CH ₃	6-CH ₃		
9.006		2-CH ₃	5-CH ₃		
9.007		3-CH ₃	5-CH ₃		
9.008		2-CH ₃	3-CH ₃		
9.009		3-CH ₃	4-CH ₃		
9.010	2-CH ₃	4-CH ₃	6-CH ₃		
9.011	2-CH ₃	4-CH ₃	5-CH ₃		
9.012		2-C1			
9.013		4-C1			
9.014		2-C1	4-Cl		
9.015		2-C1	6-C1		
9.016		2-C1	6-F		
9.017		2-CH ₃	4-C1		
9.018		2-CH ₃	4-F		
9.019		2-Cl	4-CH ₃		
9.020		2-Cl	6-CH ₃		
9.021		2-F	4-F		
9.022		2-F	6-F		
9.023		2-CH ₃	4-O-CH ₃		
9.024		2-CH ₃	6-O-CH ₃		
9.025		2-Cl	4-O-CH ₃		

- 53 -

Cmpd R ₄	R ₄	R ₄	R ₅	
9.026	2-Cl	6-O-CH ₃		
9.027	3-OCH ₃	4-OCH ₃		
9.028	2-OCH ₃	5-OCH ₃		
9.029	2-OCH ₃	4-OCH ₃		
9.030	2-OCH ₃	6-OCH ₃		
9.031	2-CF ₃	6-CF ₃		
9.032	2-CF ₃	4-CF ₃		
9.033	3-CF ₃	5-CF ₃		
9.034	2-Cl	4-CF ₃		
9.035	2-Cl	6-CF ₃		
9.036	2-NO ₂	4-NO ₂		
9.037	2-C1	4-NO ₂		
9.038	2-CH ₃	4-NO ₂		
9.039	2-O-CH ₃	4-NO ₂		
9.040	2-F	6-NO ₂		
9.041	2-Cl	6-NO ₂		
9.042	2-CH ₃	6-NO ₂		
9.043	2-O-CH ₃	6-NO ₂		
9.044	2-F	4-NO ₂		
9.045	2-CH ₃	$4-N(C_2H_5)_2$		
9.046	2-Cl	4-SO ₂ -CH ₃		
9.047	2-Cl	4-SO-CH ₃		
9.048	2-C1	4-S-CH ₃		
9.049	2-C1	6-SO ₂ -CH ₃		
9.050	2-C1	6-SO-CH ₃		
9.051	2-C1	6-S-CH ₃		
9.052	2-CH ₃	4-SO ₂ -CH ₃		
9.053	2-CH ₃	4-SO-CH ₃		
9.054	2-CH ₃	4-S-CH ₃		
9.055	2-CH ₃	6-SO ₂ -CH ₃		
9.056	2-CH ₃	6-SO-CH ₃		
9.057	2-CH ₃	6-S-CH ₃		
9.058	2-O-CH ₃	6-SO ₂ -CH ₃		
9.059	2-O-CH ₃	6-SO-CH ₃		
9.060	2-O-CH ₃	6-S-CH ₃		
9.061	2-O-CH ₃	4-SO ₂ -CH ₃		
•				

- 54 -

Cmpd	R ₄ R ₄		R ₄	R ₅
9.062		2-O-CH ₃	4-SO-CH ₃	
9.063		2-O-CH ₃	4-S-CH ₃	
9.064		2-CH ₃	$6-N(C_2H_5)_2$	
9.065		2-Cl	6-N(CH ₃) ₂	
9.066		2-Cl	4-N(CH ₃) ₂	
9.067		2-Cl	4-CO ₂ CH ₃	
9.068		2-CH ₃	6-CO ₂ C ₂ H ₅	
9.069		2-CH ₃	4-CO ₂ C ₂ H ₅	·
9.070		2-CH ₃	4-CN	
9.071		2-CH ₃	6-CN	
9.072		2-C1	4-CN	
9.073		2-Cl	6-CN	•
9.074		2-Cl	4-CO-CH ₃	
9.075		2-O-CHF ₂	4-O-CHF ₂	
9.076		2-CH ₃	4-O-CHF ₂	
9.077		2-C1	4-O-CF ₃	
9.078		2-O-CF ₃	4-O-CH ₃	
9.079		2-O-CHF ₂	4-C1	
9.080		2-O-CHF ₂	6-CH ₃	
9.081		2-O-CHF ₂	6-C1	
9.082	2-O-CHF ₂	4-CH ₃	6-CH ₃	
9.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃	
9.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇	
9.085	2-CH ₃	4-O-CH ₃	6-CH ₃	
9.086	2-C1	4-CF ₃	6-C1	
9.087	2-Cl	4-CF ₃	6-F	
9.088	2-C1	4-NO ₂	6-C1	
9.089	2-C1	4-C1	6-C1	
9.090	2-F	4-F	6-F	
9.091	2-CH ₃	4-NO ₂	6-CH ₃	
9.092	2-Cl	4-C1	6-CH ₃	
9.093	2-Cl	4-O-CH ₃	6-C1	
9.094	2-Cl	4-Cl	6-O-CH ₃	
9.095	2-F	4-O-CH ₃	6-F	
9.096	2-O-CH ₃	4-CH ₃	6-O-CH₃	
9.097	2-O-CH ₃	4-O-CH ₃	6-CH ₃	
9.098	2-O-CH ₃	4-O-CH ₃	6-O-CH ₃	
2.070				

- 55 -

Cmpd	R ₄	R ₄	R ₄	R ₅
9.099	2-O-CH ₃	4-CO-O-CH ₃	6-O-CH ₃	
9.100	2-CH ₃	4-C1	6-CH ₃	
9.101	2-CH ₃	4-F	6-CH ₃	
9.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
9.103	2-F	4-C1	5-O-i-C ₃ H ₇	
9.104	2-C1	4-C1	5-O-CH ₃	
9.105		4-Cl	5-O-CH ₃	
9.106	2-F	4-C1	5-CO-O-CH ₃	
9.107	2-F	4-Cl	5-CO-O-C ₂ H ₅	
9.108		4-C1	5-CO-O-CH ₃	
9.109	2-Cl	4-Cl	5-CO-O-i-C ₃ H ₇	
9.110				4 - 0
9.111				4 - 0 — C1
9.112				4 - 0 - F
9.113				4 - 0 - CF ₃
9.114	2-CH ₃			4-0-
9.115				4 - S - 🔷
9.116				4 - s — CI
9.117				4 - CH ₂
9.118				4 -CH ₂ -CI
9.119				4 CH ₂ - F

WO 96/11574			2199636	PCT/EP95/0393
Cmpd	R ₄	R_4	- 56 - R ₄	R ₅
9.120				4-CH ₂ - CF ₃
9.121				4 - N CHO
9.122	2-F	4-Cl	5-O-CH ₂ -CH=CH ₂	

5-O-CH₂-C=CH

9.123 2-F 4-Cl

- 57 -

Table 10: Compounds of formula Im

Cmpd	R ₄	R ₄	R ₄	R ₅	
10.001		Н			
10.002		2-CH ₃			
10.003		4-CH ₃			•
10.004		2-CH ₃	4-CH ₃		
10.005		2-CH ₃	6-CH ₃		
10.006		2-CH ₃	5-CH ₃		
10.007		3-CH ₃	5-CH ₃		
10.008		2-CH ₃	3-CH ₃		
10.009		3-CH ₃	4-CH ₃		
10.010	2-CH ₃	4-CH ₃	6-CH ₃		
10.011	2-CH ₃	4-CH ₃	5-CH ₃		
10.012		2-C1			
10.013		4-C1			
10.014		2-C1	4-C1		
10.015		2-C1	6-C1		
10.016		2-C1	6-F		
10.017		2-CH ₃	4-C1		
10.018		2-CH ₃	4-F		
10.019		2-Cl	4-CH ₃		
10.020		2-C1	6-CH ₃		
10.021		2-F	4-F		
10.022		2-F	6-F		
10.023		2-CH ₃	4-O-CH ₃		
10.024		2-CH ₃	6-O-CH ₃		
10.025		2-C1	4-O-CH ₃		
10.026		2-C1	6-O-CH ₃		

-58-

Cmpd	R ₄	R ₄	R ₄	R ₅
10.027		3-OCH₃	4-OCH ₃	
10.028		2-OCH ₃	5-OCH ₃	
10.029		2-OCH ₃	4-OCH ₃	
10.030		2-OCH ₃	6-OCH ₃	
10.031		2-CF ₃	6-CF ₃	
10.032		2-CF ₃	4-CF ₃	
10.033		3-CF ₃	5-CF ₃	
10.034		2-Cl	4-CF ₃	
10.035		2-Cl	6-CF ₃	
10.036		2-NO ₂	4-NO ₂	
10.037		2-Cl	4-NO ₂	
10.038		2-CH ₃	4-NO ₂	
10.039		2-O-CH ₃	4-NO ₂	
10.040		2-F	6-NO ₂	
10.041		2-Cl	6-NO ₂	
10.042		2-CH ₃	6-NO ₂	
10.043		2-O-CH ₃	6-NO ₂	
10.044		2-F	4-NO ₂	
10.045		2-CH ₃	$4-N(C_2H_5)_2$	
10.046		2-C1	4-SO ₂ -CH ₃	
10.047		2-C1	4-SO-CH ₃	
10.048		2-C1	4-S-CH ₃	
10.049		2-C1	6-SO ₂ -CH ₃	
10.050		2-Cl	6-SO-CH ₃	
10.051		2-C1	6-S-CH ₃	
10.052		2-CH ₃	4-SO ₂ -CH ₃	
10.053		2-CH ₃	4-SO-CH ₃	
10.054		2-CH ₃	4-S-CH ₃	
10.055		2-CH ₃	6-SO ₂ -CH ₃	
10.056		2-CH ₃	6-SO-CH ₃	
10.057		2-CH ₃	6-S-CH ₃	
10.058		2-O-CH ₃	6-SO ₂ -CH ₃	
10.059		2-O-CH ₃	6-SO-CH ₃	
10.060		2-O-CH ₃	6-S-CH ₃	
10.061		2-O-CH ₃	4-SO ₂ -CH ₃	
10.062		2-O-CH ₃	4-SO-CH ₃	

-59-

Cmpd	R ₄	R ₄	R ₄	R ₅	
10.063		2-O-CH ₃	4-S-CH ₃		
10.064		2-CH ₃	6-N(C ₂ H ₅) ₂		
10.065		2-C1	6-N(CH ₃) ₂		
10.066		2-C1	4-N(CH ₃) ₂		
10.067		2-CI	4-CO ₂ CH ₃		
10.068		2-CH ₃	6-CO ₂ C ₂ H ₅		
10.069		2-CH ₃	4-CO ₂ C ₂ H ₅		
10.070		2-CH ₃	4-CN		
10.071		2-CH ₃	6-CN		
10.072		2-C1	4-CN		
10.073		2-C1	6-CN		
10.074		2-Cl	4-CO-CH ₃		
10.075		2-O-CHF ₂	4-0-CHF ₂		
10.076		2-CH ₃	4-O-CHF ₂		
10.077		2-Cl	4-O-CF ₃		
10.078		2-O-CF ₃	4-O-CH ₃		
10.079		2-O-CHF ₂	4-C1		
10.080		2-O-CHF ₂	6-CH ₃	•	
10.081		2-O-CHF ₂	6-CI		
10.082	2-O-CHF ₂	4-CH ₃	6-CH ₃		
10.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃		
10.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇		
10.085	2-CH ₃	4-O-CH ₃	6-CH ₃		
10.086	2-C1	4-CF ₃	6-CI		
10.087	2-CI	4-CF ₃	6-F		
10.088	2-C1	4-NO ₂	6-CI		
10.089	2-C1	4-Cl	6-CI		
10.090	2-F	4-F	6-F		
10.091	2-CH ₃	4-NO ₂	6-CH ₃		
10.092	2-Cl	4-C1	6-CH ₃		
10.093	2-C1	4-O-CH ₃	6-C1		
10.094	2-C1	4-CI	6-O-CH₃		
10.095	2-F	4-O-CH ₃	6-F		
10.096	2-O-CH ₃	4-CH ₃	6-O-CH₃		
10.097	2-O-CH ₃	4-O-CH ₃	6-CH ₃		
10.098	2-O-CH ₃	4-O-CH ₃	6-O-CH ₃		

- 60 -

Cmpd	R ₄	R ₄	R ₄	R ₅
10.099	2-O-CH ₃	4-CO-O-CH₃	6-O-CH ₃	
10.100	2-CH ₃	4-C1	6-CH ₃	
10.101	2-CH ₃	4-F	6-CH ₃	
10.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
10.103	2-F	4-Cl	5-O-i-C ₃ H ₇	
10.104	2-Cl	4-Cl	5-O-CH ₃	
10.105		4-C1	5-O-CH ₃	
10.106	2-F	4-Cl	5-CO-O-CH ₃	
10.107	2-F	4-Cl	5-CO-O-C ₂ H ₅	
10.108		4-Cl	5-CO-O-CH ₃	
10.109	2-C1	4-C1	5-CO-O-i-C ₃ H ₇	
10.110				4 - 0 -
10.111				4 - 0 — CI
10.112				4 - O - F
10.113				4 O - CF ₃
10.114	2-CH ₃			4-0-
10.115				4 - S - 📞
10.116				4 - S — CI
10.117				4 - CH ₂
10.118				4 - CH ₂ - CI
10.119				4 ·CH ₂ - F

- 61 -

Cmpd	R ₄	R ₄	R ₄	R ₅
10.120				4 CH ₂ - CF ₃
10.121				4 - N - CHO
10.122	2-F	4-Cl	5-O-CH ₂ -CH=CH ₂	
10.123	2-F	4-Cl	5-O-CH ₂ -C≡CH	
10.124	2-Br			
10.125	2-CF ₃			
10.126	2-OCH ₃			•
				Cį
10.127	2-CH ₃			4-0-C1
10.128	2-CH ₃			4-0-CF ₃
10.129	2-CH ₃			4-0-C1
10.130	2-CH ₃		6-CH₃	4-0-
10.131	2-CH ₃		6-CH ₃	4 O - CF ₃
10.132	2-CH ₃		6-CH₃	4-0-CI
10.133	2-CH ₃		6-CH ₃	4-0-C1
10.134	2-CH ₃	4-Br	6-CH₃	CI
10.135	2-CH ₃	6-C ₂ H ₅		
10.136	$2-C_2H_5$	6-C ₂ H ₅		
10.137	2-CH ₃	4-OC ₂ H ₅	6-CH ₃	
10.138	2-CH ₃	4-O-i-C ₃ H ₇	6-CH ₃	
10.139		4-O-n-C ₃ H ₇	6-CH ₃	
10.140	2-CH ₃	4-O-n-C ₁₀ H ₂₁	6-CH ₃	

- 62 -

Cmpd	R ₄	R ₄	R ₄	R ₅
10.141	2-CH ₃	4-O-n-C ₃ H ₇		
10.142	2-CH ₃	4-O-(CH ₂) ₂ OCH ₃		
10.143	2-CH ₃	4-O-(CH ₂) ₂ OCH ₃	6-CH ₃	
10.144	2-CH ₃	4-O-(CH ₂) ₂ OCH ₃		
10.145	2-CH ₃	4-O-n-C ₆ H ₁₃	6-CH ₃	
10.146	2-CH ₃			4-0-
10.147	2-CH ₃		6-CH ₃	4 -o — N
10.148	2-CH ₃			4 -O -CF ₃
10.149	2-CH ₃		6-CH ₃	4 -O -CF ₃
10.150	2-CH ₃			4-0-N-CF ₃
				ď
10.151	2-CH ₃		6-CH ₃	4-0-CF ₃
				cí N—
10.152	2-CH ₃	•		5-0-CF ₃
				CÍ N
10.153	2-CH ₃			5-0-CF ₃
10.154	2-CH ₃		6-CH ₃	4 - s —
10.155	2-CH ₃		6-CH ₃	4-s-(a
10.156	2-C ₂ H ₅		6-CH₃	4 - S —
10.157 10.158	2-C ₂ H ₅ 2-C ₂ H ₅	4-C ₂ H ₅	6-C ₂ H ₅	_

- 63 -

10.159 2-CH(CH₃)₂

Table 11: Compounds of formula In

Cmpd	R ₄	R ₄	R ₄	R ₅	
11.001		Н			
11.002		2-CH ₃			
11.003		4-CH ₃			
11.004		2-CH ₃	4-CH ₃		
11.005		2-CH ₃	6-CH ₃		
11.006		2-CH ₃	5-CH ₃		
11.007		3-CH ₃	5-CH ₃		:
11.008		2-CH ₃	3-CH ₃		
11.009		3-CH ₃	4-CH ₃		
11.010	2-CH ₃	4-CH ₃	6-CH ₃		
11.011	2-CH ₃	4-CH ₃	5-CH ₃		
11.012		2-C1			
11.013		4-C1			
11.014		2-C1	4-CI		
11.015		2-C1	6-C1		
11.016		2-Cl	6-F		
11.017		2-CH ₃	4-C1		
11.018		2-CH ₃	4-F		
11.019		2-C1	4-CH ₃		
11.020		2-C1	6-CH ₃		
11.021		2-F	4-F		
11.022		2-F	6-F		
11.023		2-CH ₃	4-O-CH ₃		
11.024		2-CH ₃	6-O-CH ₃		
11.025		2-C1	4-O-CH ₃		
11.023		2-U	4-U-LH ₃		

- 64 -

Cmpd	R ₄	R ₄	R ₄	R ₅
11.026	<u>-</u>	2-Cl	6-O-CH₃	,
11.027		3-OCH ₃	4-OCH ₃	
11.028		2-OCH ₃	5-OCH ₃	
11.029		2-OCH ₃	4-OCH ₃	
11.030		2-OCH ₃	6-OCH ₃	
11.031		2-CF ₃	6-CF ₃	
11.032		2-CF ₃	4-CF ₃	
11.033		3-CF ₃	5-CF ₃	
11.034		2-C1	4-CF ₃	
11.035		2-Cl	6-CF ₃	
11.036		2-NO ₂	4-NO ₂	
11.037		2-C1	4-NO ₂	
11.038		2-CH ₃	4-NO ₂	
11.039		2-O-CH ₃	4-NO ₂	
11.040		2-F	6-NO ₂	
11.041		2-C1	6-NO ₂	
11.042		2-CH ₃	6-NO ₂	
11.043		2-O-CH ₃	6-NO ₂	
11.044		2-F	4-NO ₂	
11.045		2-CH ₃	$4-N(C_2H_5)_2$	
11.046		2-C1	4-SO ₂ -CH ₃	
11.047		2-C1	4-SO-CH ₃	
11.048		2-C1	4-S-CH ₃	
11.049		2-Cl	6-SO ₂ -CH ₃	
11.050		2-C1	6-SO-CH ₃	
11.051		2-Cl	6-S-CH ₃	
11.052		2-CH ₃	4-SO ₂ -CH ₃	
11.053		2-CH ₃	4-SO-CH ₃	
11.054		2-CH ₃	4-S-CH ₃	
11.055		2-CH ₃	6-SO ₂ -CH ₃	
11.056		2-CH ₃	6-SO-CH₃	
11.057		2-CH ₃	6-S-CH ₃	
11.058		2-O-CH ₃	6-SO ₂ -CH ₃	
11.059		2-O-CH ₃	6-SO-CH ₃	
11.060		2-O-CH ₃	6-S-CH ₃	
11.061		2-O-CH ₃	4-SO ₂ -CH ₃	
		-		

- 65 -

Cmpd	R ₄	R ₄	R ₄	R ₅
11.062		2-O-CH ₃	4-SO-CH ₃	
11.063		2-O-CH ₃	4-S-CH ₃	
11.064		2-CH ₃	6-N(C ₂ H ₅) ₂	
11.065		2-Cl	6-N(CH ₃) ₂	
11.066		2-Cl	4-N(CH ₃) ₂	
11.067		2-Cl	4-CO ₂ CH ₃	
11.068		2-CH ₃	6-CO ₂ C ₂ H ₅	
11.069		2-CH ₃	4-CO ₂ C ₂ H ₅	
11.070		2-CH ₃	4-CN	
11.071		2-CH ₃	6-CN	
11.072		2-Cl	4-CN	
11.073		2-Cl	6-CN	
11.074		2-CI	4-CO-CH ₃	
11.075		2-O-CHF ₂	4-O-CHF ₂	
11.076		2-CH ₃	4-O-CHF ₂	
11.077		2-Cl	4-O-CF ₃	
11.078		2-O-CF ₃	4-O-CH ₃	
11.079		2-O-CHF ₂	4-C1	
11.080		2-O-CHF ₂	6-CH ₃	,
11.081		2-O-CHF ₂	6-C1	
11.082	2-O-CHF ₂	4-CH ₃	6-CH ₃	
11.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃	
11.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇	
11.085	2-CH ₃	4-O-CH ₃	6-CH ₃	
11.086	2-CI	4-CF ₃	6-C1	
11.087	2-C1	4-CF ₃	6-F	
11.088	2-C1	4-NO ₂	6-C1	
11.089	2-C1	4-Cl	6-C1	
11.090	2-F	4-F	6-F	
11.091	2-CH ₃	4-NO ₂	6-CH ₃	
11.092	2-C1	4-CI	6-CH ₃	
11.093	2-Cl	4-O-CH ₃	6-C1	
11.094	2-Cl	4-C1	6-O-CH ₃	
11.095	2-F	4-O-CH ₃	6-F	
11.096	2-O-CH ₃	4-CH ₃	6-O-CH ₃	
11.097	2-O-CH ₃	4-O-CH ₃	6-CH ₃	

- 66 -

Cmpd	R ₄	R ₄	R ₄	R ₅
11.098	2-O-CH ₃	4-O-CH ₃	6-O-CH ₃	
11.099	2-O-CH ₃	4-CO-O-CH ₃	6-O-CH ₃	
11.100	2-CH ₃	4-C1	6-CH ₃	
11.101	2-CH ₃	4-F	6-CH ₃	
11.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
11.103	2-F	4-Cl	5-O-i-C ₃ H ₇	
11.104	2-C1	4-C1	5-O-CH ₃	
11.105		4-CI	5-O-CH ₃	
11.106	2-F	4-C1	5-CO-O-CH ₃	
11.107	2-F	4-C1	5-CO-O-C ₂ H ₅	
11.108		4-C1	5-CO-O-CH ₃	
11.109	2-C1	4-C1	5-CO-O-i-C ₃ H ₇	
11.110				4 - 0 -
11.111				4 - 0 — CI
11.112				4 - 0 — F
11.113				4-O-CF ₃
11.114	2-CH ₃			4-0-
11.115				4 - S —
11.116				4 - S — CI
11.117				4 - CH ₂ —
11.118				4 CH ₂ - CI
11.119				4 CH ₂ — F

- 67 -

Cmpd	R ₄	R ₄	R ₄	R ₅
11.120				4-CH ₂ -CF ₃
11.121				4 - N-(N-(N-(N-(N-(N-(N-(N-(N-(N-(N-(N-(N-(N
11.122	2-F	4-Cl	5-O-CH ₂ -CH=CH ₂	
11.123	2-F	4-Cl	5-O-CH ₂ -C=CH	
11.124	2-Br		_	
11.125	2-CF ₃			
11.126	2-OCH ₃			
				CI
11.127	2-CH ₃			4 -0 — CI
11.128	2-CH ₃			4-0-CF ₃
11.129	2-CH ₃			4 -0 — C1
11.130	2-CH ₃		6-CH ₃	4 -0 -
11.131	2-CH ₃		6-CH ₃	40-CF3
11.132	2-CH ₃		6-CH₃	4 -0 — C1
11.133	2-CH ₃		6-CH ₃	4-0-CI
11.134	2-CH ₃	4-Br	6-CH ₃	ĊI
11.135	2-CH ₃	6-C ₂ H ₅		
11.136	$2-C_2H_5$			
11.137		4-OC ₂ H ₅	6-CH ₃	
11.138		4-O-i-C ₃ H ₇	6-CH ₃	
11.139	=	4-O-n-C ₃ H ₇	6-CH ₃	
11.140	2-CH ₃	4-O-n-C ₁₀ H ₂₁	6-CH ₃	

- 68 -

Cmpd	R ₄	R ₄	R ₄	R ₅
11.141	2-CH ₃	4-O-n-C ₃ H ₇		
11.142	2-CH ₃	4-O-(CH ₂) ₂ OCH ₃		
	_	4-O-(CH ₂) ₂ OCH ₃	6-CH ₃	
		4-O-(CH ₂) ₂ OCH ₃		
11.145	2-CH ₃	4-O-n-C ₆ H ₁₃	6-CH ₃	N
11.146	2-CH ₃			4-0-
11.147	2-CH ₃		6-CH ₃	4-0-
11.148	2-CH ₃			4 -0 - CF ₃
11.149	2-CH ₃		6-CH ₃	4 .0 — CF ₃
11.150	2-CH ₃			4-0-KP3
11.151	2-CH ₃		6-CH ₃	Cí 4-0
11.152	2-CH ₃			5-0-X-CF ₃
11.153	2-CH ₃			5 -o — CF ₃
11.154	2-CH ₃		6-CH ₃	4 - s —
11.155	2-CH ₃		6-CH ₃	4 - s — C1
11.156	2-C ₂ H ₅		6-CH ₃	4 - s —

- 69 -

Table 12: Compounds of formula Io

$$\begin{array}{c|c}
CH_3 & O \\
CH_2 & C & N
\end{array}$$

$$\begin{array}{c|c}
(R_4)_n \\
(R_5)_m
\end{array}$$
(Io)

Cpmd	R ₄	R ₄	R ₄	R ₅	
12.001		Н			· · · · · · · · · · · · · · · · · · ·
12.002		2-CH ₃			
12.003		4-CH ₃			
12.004		2-CH ₃	4-CH ₃		
12.005		2-CH ₃	6-CH ₃		
12.006		2-CH ₃	5-CH ₃		
12.007		3-CH ₃	5-CH ₃		
12.008		2-CH ₃	3-CH ₃		· •
12.009		3-CH ₃	4-CH ₃		*see
12.010	2-CH ₃	4-CH ₃	6-CH ₃		
12.011	2-CH ₃	4-CH ₃	5-CH ₃		
12.012		2-C1			
12.013		4-C1			
12.014		2-C1	4-C1		
12.015		2-C1	6-Cl		
12.016		2-Cl	6-F		
12.017		2-CH ₃	4-C1		
12.018		2-CH ₃	4-F		
12.019		2-Cl	4-CH ₃		
12.020		2-C1	6-CH ₃		
12.021		2-F	4-F		
12.022		2-F	6-F		
12.023		2-CH ₃	4-O-CH ₃		
12.024		2-CH ₃	6-O-CH ₃		
12.025		2-C1	4-O-CH ₃		
12.026		2-C1	6-O-CH₃		

- 70 -

Cmpd R ₄	R ₄	R ₄	R ₅
12.027	3-OCH ₃	4-OCH ₃	
12.028	2-OCH ₃	5-OCH ₃	
12.029	2-OCH ₃	4-OCH ₃	
12.030	2-OCH ₃	6-OCH ₃	
12.031	2-CF ₃	6-CF ₃	
12.032	2-CF ₃	4-CF ₃	
12.033	3-CF ₃	5-CF ₃	
12.034	2-Cl	4-CF ₃	
12.035	2-Cl	6-CF ₃	
12.036	2-NO ₂	4-NO ₂	
12.037	2-Cl	4-NO ₂	
12.038	2-CH ₃	4-NO ₂	
12.039	2-O-CH ₃	4-NO ₂	
12.040	2-F	6-NO ₂	
12.041	2-Cl	6-NO ₂	
12.042	2-CH ₃	6-NO ₂	
12.043	2-O-CH ₃	6-NO ₂	
12.044	2-F	4-NO ₂	
12.045	2-CH ₃	$4-N(C_2H_5)_2$	
12.046	2-Cl	4-SO ₂ -CH ₃	
12.047	2-C1	4-SO-CH ₃	
12.048	2-Cl	4-S-CH ₃	
12.049	2-Cl	6-SO ₂ -CH ₃	
12.050	2-Cl	6-SO-CH ₃	
12.051	2-Cl	6-S-CH ₃	
12.052	2-CH ₃	4-SO ₂ -CH ₃	
12.053	2-CH ₃	4-SO-CH ₃	
12.054	2-CH ₃	4-S-CH ₃	
12.055	2-CH ₃	6-SO ₂ -CH ₃	
12.056	2-CH ₃	6-SO-CH₃	
12.057	2-CH ₃	6-S-CH ₃	
12.058	2-O-CH ₃	6-SO ₂ -CH ₃	
12.059	2-O-CH ₃	6-SO-CH ₃	
12.060	2-O-CH ₃	6-S-CH ₃	
12.061	2-O-CH ₃	4-SO ₂ -CH ₃	
12.062	2-O-CH ₃	4-SO-CH ₃	

- 71 -

Cmpd	R ₄	R ₄	R ₄	R ₅
12.063		2-O-CH ₃	4-S-CH ₃	······································
12.064		2-CH ₃	$6-N(C_2H_5)_2$	
12.065		2-C1	$6-N(CH_3)_2$	
12.066		2-C1	$4-N(CH_3)_2$	
12.067		2-C1	4-CO ₂ CH ₃	
12.068		2-CH ₃	6-CO ₂ C ₂ H ₅	·
12.069		2-CH ₃	4-CO ₂ C ₂ H ₅	
12.070		2-CH ₃	4-CN	
12.071		2-CH ₃	6-CN	
12.072		2-Cl	4-CN	
12.073		2-C1	6-CN	
12.074		2-C1	4-CO-CH ₃	
12.075		2-O-CHF ₂	4-O-CHF ₂	
12.076		2-CH ₃	4-0-CHF ₂	
12.077		2-C1	4-O-CF ₃	•
12.078		2-O-CF ₃	4-O-CH ₃	
12.079		2-O-CHF ₂	4-CI	
12.080		2-O-CHF ₂	6-CH ₃	
12.081		2-O-CHF ₂	6-CI	
12.082	2-O-CHF ₂	4-CH ₃	6-CH ₃	
12.083	2-CH ₃	4-t-C ₄ H ₉	6-CH ₃	
12.084	2-i-C ₃ H ₇	4-i-C ₃ H ₇	6-i-C ₃ H ₇	
12.085	2-CH ₃	4-O-CH ₃	6-CH ₃	
12.086	2-C1	4-CF ₃	6-CI	
12.087	2-Cl	4-CF ₃	6-F	
12.088	2-Cl	4-NO ₂	6-C1	
12.089	2-Cl	4-Cl	6-CI	
12.090	2-F	4-F	6-F	
12.091	2-CH ₃	4-NO ₂	6-CH ₃	
12.092	2-C1	4-C1	6-CH ₃	
12.093	2-Cl	4-O-CH ₃	6-CI	
12.094	2-Cl	4-Cl	6-O-CH ₃	
12.095	2-F	4-O-CH ₃	6-F	
12.096	2-O-CH ₃	4-CH ₃	6-O-CH ₃	
12.097	2-O-CH ₃	4-O-CH ₃	6-CH ₃	

Cmpd	R ₄	R ₄	R ₄	R ₅
12.098	2-O-CH ₃	4-O-CH ₃	6-O-CH₃	
12.099	2-O-CH ₃	4-CO-O-CH ₃	6-O-CH ₃	
12.100	2-CH ₃	4-C1	6-CH ₃	
12.101	2-CH ₃	4-F	6-CH ₃	
12.102	2-CH ₃	4-CH ₃	6-O-CH ₃	
12.103	2-F	4-C1	5-O-i-C ₃ H ₇	
12.104	2-C1	4-C1	5-O-CH ₃	
12.105		4-C1	5-O-CH ₃	
12.106	2-F	4-CI	5-CO-O-CH ₃	
12.107	2-F	4-C1	5-CO-O-C ₂ H ₅	
12.108		4-CI	5-CO-O-CH ₃	
12.109	2-C1	4-C1	5-CO-O-i-C ₃ H ₇	
12.110				4 - 0
12.111				4 - 0 — CI
12.112				4 - 0 — F
12.113				4-O-CF ₃
12.114	2-CH ₃			4-0-
12.115				4 - 8 -
12.116				4 - S — CI
12.117				4 - CH ₂ -
12.118				4-CH ₂ -CI
12.119				4-CH ₂ — F

- WO 96/11574

PCT/EP95/03935

- 73 -

Cmpd	R ₄	R ₄	R ₄	R ₅
12.120				4-CH ₂ - CF ₃
12.121				4 - N-(CHO)
12.122	2-F	4-Cl	5-O-CH ₂ -CH=CH ₂	
12.123	2-F	4-Cl	5-O-CH ₂ -C=CH	

2199636

- 74 -

Table 13: Compounds of formula Ip

$$\bigcap_{N} \bigvee_{O} \bigcap_{(P_{A})}$$

(Ip)

Cmpd	R ₄	
13.001	Н 171-172	
13.002	2-CH ₃	
13.003	2-OCH ₃	
13.004	4-C1	
13.005	4-F	
13.006	4-NO ₂	
13.007	6-NO ₂	
13.008	7-NO ₂	

- 75 -

Table 14: Compounds of formula Iq

$$CH_3$$
 N
 CH_3
 N
 (Iq)

Cmpd	R ₄	
14.001	Н	
14.002	2-CH ₃	
14.003	2-OCH ₃	
14.004	4-Cl	
14.005	4-F	
14.006	4-NO ₂	
14.007	6-NO ₂	
14.008	7-NO ₂	.;

- 76 -

Table 15: Compounds of formula Ir

$$\begin{array}{c|c} A & C-G & CH_3 \\ \hline & N & CH_3 \\ \hline & O & CH_3 \end{array}$$
 (Ir)

Cmpd	G	А В	
15.001	Н	-(CH ₂) ₄ -	
15.002	-C-CH ₂ -C(CH ₃) ₃	-(CH ₂) ₄ -	
15.003	-CO-C(CH ₃) ₃	-(CH ₂) ₄ -	
15.004	-COOCH(CH ₃) ₂	-(CH ₂) ₄ -	
15.005	-COOC ₂ H ₅	-(CH ₂) ₄	
15.006	H	H C ₆ H ₅	
15.007	-C-N(CH ₃) ₂ S OCH ₃	-(CH ₂) ₄ -	
15.008	-PCH ₃	-(CH ₂) ₄ -	
15.009	OCH ₃ C ₂ H ₅	-(CH ₂) ₄ -	
15.010	CH ₃ -CO-C-C ₂ H ₅ CH ₃	-(CH ₂) ₄ -	
15.011	-CO-CH(CH ₃) ₂	-(CH ₂) ₄ -	

- 77 -

Cmpd	G	A B
15.012	OC ₂ H ₅ -P SCH-C ₂ H ₅ O CH ₃	-(CH ₂) ₄ -
15.013	CH ₃ -COCCH₂CH₂CI CH ₃	-(CH ₂) ₄ -
15.014	н	-CH ₂ CF-CH ₂ - CH ₃
15.015	-COSC ₂ H ₅	-(CH ₂) ₄ -
15.016	-CON(CH ₃) ₂ CH ₃	-(CH ₂) ₄ -
15.017	-COC-C ₂ H ₅	-CH ₂ CF ₂ CH ₂ -
15.018	-COCH ₃	-CH ₂ CF-CH ₂ - CH ₃
15.019	-COC(CH ₃) ₃	-CH ₂ CF-CH ₂ - CH ₃
15.020	Н	-(CH ₂) ₃ -
15.021	-COOCH-CH₃ CH₂OC₂H₅ ÇH₃	-CH ₂ CF-CH ₂ - CH ₃
15.022	-COCH ₂ C-CH ₃ CH ₃	-CH ₂ CF-CH ₂ CH ₃

- 78 -

Cmpd	G	А В
15.023	CH ₃ -COCH ₂ C-CH ₃ CH ₃	-(CH ₂) ₃ -
15.024	-COC-CH CH ₃ CH ₃	-(CH ₂) ₄ -
15.025	-CO-C(CH ₃) ₃	-(CH ₂) ₃ -
15.026	-CO-C(CH ₃) ₃	-CH ₂ CF ₂ CH ₂ -
15.027	-COCH ₃	H C ₆ H ₅
15.028	-COCH ₂ C(CH ₃) ₃	H C ₆ H ₅
15.029	-COC(CH ₃) ₃	H C ₆ H ₅
		Λ
15.030	Н	-CH ₂
15.031	H	-CH ₂ CF ₂ CH ₂ -
15.032	CH ₃ -COCH ₂ C-CH ₃ CH ₃	-CH ₂ CF ₂ CH ₂ -
15.033	-COCH ₃	-CH ₂ CF ₂ CH ₂ -
15.034	-COOCH-CH ₂ O CH ₃ CH ₃	-CH ₂ CF ₂ CH ₂ -
15.035	-COC CH CH ₃ CH ₃ -CH CH ₃ CH ₃	-CH ₂ CF ₂ CH ₂ -
15.036	-SO ₂ CH ₃	-(CH ₂) ₄ -
15.037	Н	н — н

- 79 -

Cmpd	G	A B
15.038	-C-O-CH-CH ₂ -OC ₂ H ₅ O CH ₃	
15.039	-C-C(CH ₃) ₂ -CH(CH ₃) ₂ O	
15.040	-C-C(CH ₃) ₂ -C ₂ H ₅	
15.041	-C-CH₃ O	
15.042	-C-C(CH ₃) ₃ O	
15.043	-C-C(CH ₃) ₂ -CH(CH ₃) ₂ O	н Н
15.044	-C-O-CH-CH ₂ -OC ₂ H ₅ O CH ₃	н н
15.045	-C-C(CH ₃) ₂ -C ₂ H ₅	н — н
15.046	Н	-(CH ₂) ₅ -
15.047	Н	-CH ₂ -C(CH ₃) ₂ -CH ₂ -
15.048	-C-CH ₂ -C(CH ₃) ₃	-CH ₂ -C(CH ₃) ₂ -CH ₂ -
15.049	-C-C(CH ₃) ₂ -C ₂ H ₅	-(CH ₂) ₅ -
15.050	-C-CH₃ O	-(CH ₂) ₅ -
15.051	н	-CH(CH ₃)-CH ₂ -CH(CH ₃)-

- 80 -

Cmpd	G	A B	
15.052	Н	н СН ₃	
15.053	-C-C(CH ₃) ₃	H CH ₃	
15.054	-C-C(CH ₃) ₂ -C ₂ H ₅ O	H CH ₃	
15.055	-C-CH ₂ -C(CH ₃) ₃ O	H CH ₃	
15.056	-C-CH₃ O	H CH ₃	
15.057	-C-C(CH ₃) ₃	-CH(CH ₃)-CH ₂ -CH(CH ₃)-	
15.058	-C-CH₃ O	-CH(CH ₃)-CH ₂ -CH(CH ₃)-	
15.059	-C-C(CH ₃) ₂ -CH(CH ₃) ₂ O	-CH(CH ₃)-CH ₂ -CH(CH ₃)-	
15.060	-C-CH ₂ -C(CH ₃) ₃ O	-CH(CH ₃)-CH ₂ -CH(CH ₃)-	
15.061	-C-C(CH ₃) ₂ -C ₂ H ₅ 0	-CH(CH ₃)-CH ₂ -CH(CH ₃)-	
15.062	-C-C(CH ₃) ₃ O	-CH ₂ -C(C ₃ H ₇)-CH ₂ - CH ₃	
15.063	-C-C(CH ₃) ₂ -CH(CH ₃) ₂ O	-CH ₂ -C(C ₃ H ₇)-CH ₂ - CH ₃	
15.064	-C-C(CH ₃) ₃	-CH ₂ -CH ₂ -CH- CH ₃	

- 81 -

Cmpd	G	А В	
15.065	-C-C(CH ₃) ₂ -CH(CH ₃) ₂ O	-CH ₂ -CH ₂ -CH- CH ₃	
15.066	-C-CH₃ O	-CH ₂ -C(C ₃ H ₇)-CH ₂ - CH ₃	
15.067	– СС(СН ₃) ₃ Ö	CH ₃ CH ₃	
15.068	– ССН ₂ С(СН ₃) ₃ Ö	CH ₃ CH ₃	
15.069	Н	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -	•
15.070	– CC(CH₃)₃ Ö	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -	
15.071	– CCH₂C(CH₃)₃ Ö	-(CH ₂) ₂ CH(CH ₃)(CH ₂) ₂ -	
15.072	− CC(CH ₃) ₃ Ö	-CH ₂ C(CH ₃) ₂ CH ₂ -	٠.
15.073	- СС(СН ₃) ₃ Ö	-CH ₂ CH(CH ₃)CH ₂ CH ₂ -	
15.074	– CC(CH₃)₃ Ö	$-CH_3$ $-C_2H_5$	
15.075	– СС(СН₃)₃ Ö	-CH(CH ₃)(CH ₂) ₃ -	
15.076	– CC(CH₃)₃ Ö	-CH ₂ CH(CH ₃)CH(CH ₃)CH ₂ -	

Preferred compounds of formula IIa are listed in the following Table 16.

2199636

- 82 -

Table 16: Compounds of formula IIa:

$$\begin{array}{c|c}
X_2 \\
\hline
O-CH_2-C-O-R_{10} \\
\hline
0
\end{array}$$
(IIa),

Cmpd	X_2	R ₁₀	
16.01	Cl	-CH(CH ₃)-C ₅ H ₁₁ -n	
16.02	Cl	-CH(CH ₃)-CH ₂ OCH ₂ CH=CH ₂	
16.03	Cl	Н	
16.04	Cl	C ₄ H ₉ -n	

Preferred compounds of formula IIb and IIb₁ are listed in the following Tables 17 and 18.

Table 17: Compounds of formula IIb:

$$R_{11}$$

$$R_{12}$$

$$R_{13}$$
(IIb)

Cmpd	R ₁₄	R ₁₁	R ₁₂	R ₁₃	E
17.01	CH ₃	phenyl	2-C1	Н	CH
17.02	CH ₃	phenyl	2-Cl	4-C1	CH
17.03	CH ₃	phenyl	2-F	H	CH
17.04	CH ₃	2-chlorophenyl	2-F	Н	СН
17.05	C_2H_5	CCl ₃	2-C1	4-C1	N
17.06	CH ₃	phenyl	2-C1	4-CF ₃	N

- 83 -

Cmpd	R ₁₄	R ₁₁	R ₁₂	R ₁₃	E	
17.07	CH₃	phenyl		2-C1	4-CF ₃	N

Table 18: Compound of formula IIb₁:

$$R_{67}OOC$$
 R_{68}
 R_{12}
 R_{13}
 R_{13}
 R_{13}
 R_{13}
 R_{14}
 R_{15}

Cmpd	R ₆₈	R ₆₇	R ₆₆	R ₁₂	R ₁₃
18.01	CH ₃	CH ₃	CH ₃	2-C1	4-Cl
18.02	CH ₃	C ₂ H ₅	CH ₃	2-C1	4-C1
18.03	CH₃	C ₂ H ₅	C_2H_5	2-C1	4-CI

Table 19: Compounds of formula Is

$$\begin{array}{c|c}
A & O-G & R_4 \\
N & & & \\
N & & & \\
O & R_4
\end{array}$$
(Is)

Cmpd	R ₄	R ₄	R ₄	G	A	В
19.01	C ₂ H ₅	C ₂ H ₅	C₂H₅	— CCH ₂ C(CH ₃) ₃ Ⅱ O		-(CH ₂) ₄ -
19.02	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	CC(CH₃)₃ Ⅱ O	CH ₃	CH ₃
19.03	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	− CC(CH ₃) ₃ Ⅱ O		-(CH ₂) ₄ -

- 84 -

Cmpd	R ₄	R ₄	R ₄	G	A	В
19.04	Cl	Н	F	CC(CH ₃) ₃ II O	CH ₃	CH ₃
19.05	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	- CCH ₃ II O		-(CH ₂) ₄ -
19.06	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	-COC ₂ H ₅ Ⅱ O		-(CH ₂) ₄ -
19.07	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	−CC ₂ H ₅		-(CH ₂) ₄ -
19.08	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	— CCH(CH ₃)₂ Ⅱ O		-(CH ₂) ₄ -

The compounds of formula I are disclosed in WO 92/16510 and EP-A-0 508 126 as insecticides, acaricides and herbicides.

The quinoline derivatives falling under the scope of formula IIa and the preparation thereof are known or can be prepared according to known processes, such as disclosed, inter alia, in EP-A-0 094 349.

The 1-phenylazole-3-carboxylic acid derivatives falling under the scope of formula IIb are known and are disclosed, inter alia, in EP-A-0 268 554 and EP-A-0 174 562.

The compounds of formula IIb₁ are disclosed in WO 91/07874 as safener.

The invention also relates to a method of selectively controlling weeds in crops of cultivated plants, which comprises treating said cultivated plants, the seeds or seedlings or the crop area thereof, concurrently or separately, with a herbicidally effective amount of the herbicide of formula I and, to antagonise the herbicide, an antidotally effective amount of the safener of formula IIa, IIb or IIb₁.

Suitable cultivated plants which can be protected by the safener of formula IIa, IIb and IIb₁ against the harmful action of the aforementioned herbicides are preferably maize and cereals.

The weeds to be controlled can be monocot as well as dicot weeds.

- 85 -

Crop areas will be understood as meaning the areas already under cultivation with the cultivated plants or seeds thereof, as well as the areas intended for cropping with said cultivated plants.

Depending on the end use, a safener of formula IIa, IIb and IIb₁ can be used for pretreating seeds of the crop plants (dressing of seeds of seedlings) or it can be incorporated in the soil before or after sowing. It can, however, also be applied by itself alone or together with the herbicide postemergence. Treatment of the plant or the seeds with the safener can therefore in principle be carried out irrespective of the time of application of the herbicide. Treatment can, however, also be carried out by simultaneous application of the phytotoxic chemical and safener (e.g. as tank mixture).

The concentration of safener with respect to the herbicide will depend substantially on the mode of application. Where a field treatment is carried out either by using a tank mixture with a combination of safener and herbicide or by separate application of safener and herbicide, the ratio of safener to herbicide will usually be from 100:1 to 1:10, preferably from 20:1 to 1:1.

In field treatment it is usual to apply 0.001 to 5.0 kg/ha, preferably 0.001 to 0.5 kg/ha, of safener.

The concentration of herbicide is usually in the range from 0.001 to 2 kg/ha, but will preferably be from 0.005 to 1 kg/ha.

The compositions of this invention are suitable for all methods of application commonly used in agriculture, including preemergence application, postemergence application and seed dressing.

For seed dressing, 0.001 to 10 g of safener/kg of seeds, preferably 0.05 to 2 g of safener/kg of seeds, is usually applied. If the safener is used in liquid form shortly before sowing to effect soaking, then it is preferred to use safener solutions that contain the active ingredient in a concentration of 1 to 10 000 ppm, preferably of 100 to 1000 ppm.

For application, it is preferred to process the safeners of formula IIa, IIb or IIb₁, or mixtures of these safeners nd the herbicides tof formula I together with the assistants

conventionally employed in formulation technology to emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates or microcapsules. The formulations are prepared in known manner, conveniently by homogeneously mixing or grinding, or mixing and grinding, the active ingredients with liquid or solid formulation assistants, typically solvents or solid carriers. Surface-active compounds (surfactants) may additionally be used for preparing the formulations.

Suitable solvents may typically be: aromatic hydrocarbons, preferably the fractions containing 8 to 12 carbon atoms such as xylene mixtures or substituted naphthalenes; phthalates such as dibutyl or dioctyl phthalate; aliphatic hydrocarbons such as cyclohexane or paraffins; alcohols and glycols and their ethers and esters such as ethanol, diethylene glycol, 2-methoxyethanol or 2-ethoxyethanol; ketones such as cyclohexanone; strongly polar solvents such as N-methyl-2-pyrrolidone, dimethyl sulfoxide or dimethyl formamide; as well as vegetable oils or epoxidised vegetable oils such as epoxidised coconut oil or soybean oil; or water.

The solid carriers typically used for dusts and dispersible powders are usually natural mineral fillers such as calcite, talcum, kaolin, montmorillonite or attapulgite. To improve the physical properties it is also possible to add highly dispersed silicic acid or highly dispersed absorbent polymers. Suitable granulated adsorptive carriers are porous types, including pumice, broken brick, sepiolite or bentonite; and suitable nonsorbent carriers are materials such as calcite or sand. In addition, innumerable pregranulated materials of inorganic or organic origin may be used, especially dolomite or pulverised plant residues.

Depending on the safener, and usually also on the herbicide, suitable surface-active compounds are nonionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. Surfactants will also be understood as comprising mixtures of surfactants.

Suitable anionic surfactants may be water-soluble soaps as well as water-soluble synthetic surface-active compounds.

Suitable soaps are the alkali metal salts, alkaline earth metal salts, ammonium salts or substituted ammonium salts of higher fatty acids $(C_{10}-C_{22})$, e.g. the sodium or potassium salts of oleic or stearic acid, or of natural fatty acid mixtures which can be obtained, inter

Q)

alia from coconut oil or tallow oil. Further suitable soaps are also the fatty acid methyl taurin salts.

More often, however, so-called synthetic surfactants are used, especially fatty sulfonates, fatty sulfates, sulfonated benzimidazole derivatives or alkylarylsulfonates.

The fatty alcohol sulfonates or sulfates are usually in the form of alkali metal salts, alkaline earth metal salts, ammonium salts or substituted ammonium salts, and they contain a C_8 - C_{22} alkyl radical which also includes the alkyl moiety of acyl radicals, e.g. the sodium or calcium salt of ligninsulfonic acid, of dodecylsulfate, or of a mixture of fatty alcohol sulfates obtained from natural fatty acids. These compounds also comprise the salts of sulfated and sulfonated fatty alcohol/ethylene oxide adducts. The sulfonated benzimidazole derivatives preferably contain two sulfonic acid groups and one fatty acid radical containing 8 to 22 carbon atoms. Illustrative examples of alkylarylsulfonates are the sodium, calcium or triethanolamine salts of dodecylbenzenesulfonic acid, dibutylnaphthalenesulfonic acid, or of a condensate of naphthalenesulfonic acid and formaldehyde.

Corresponding phosphates, typically salts of the phosphoric acid ester of an adduct of p-nonylphenol with 4 to 14 mol of ethylene oxide, or phospholipids, are also suitable.

Nonionic surfactants are preferably polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols or of saturated or unsaturated fatty acids and alkylphenols, said derivatives containing 3 to 30 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenols.

Further suitable nonionic surfactants are the water-soluble polyadducts of polyethylene oxide with polypropylene glycol, ethylenediaminopolypropylene glycol and alkylpolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which polyadducts contain 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups. These compounds usually contain 1 to 5 ethylene glycol units per propylene glycol unit.

Illustrative examples of nonionic surfactants are nonylphenol polyethoxylates, polyethoxylated castor oil, polyadducts of polypropylene and polyethylene oxide, tributylphenol polyethoxylate, polyethylene glycol and octylphenol polyethoxylate.

- 88 -

Fatty acid esters of polyoxyethylene sorbitan are also suitable nonionic surfactants, typically polyoxyethylene sorbitan trioleate.

Cationic surfactants are preferably quaternary ammonium salts carrying, as N-substituent, at least one C₈-C₂₂alkyl radical and, as further substituents, unsubstituted or halogenated lower alkyl, benzyl or hydroxy-lower alkyl radicals. The salts are preferably in the form of halides, methyl sulfates or ethyl sulfates, for example stearyl trimethylammonium chloride or benzyl bis(2-chloroethyl)ethylammonium bromide.

The surfactants customarily employed in the art of formulation are described, inter alia, in "Mc Cutcheon's Detergents and Emulsifiers Annual", Mc Publishing Corp., Glen Rock, New Jersey, 1988, H. Stache, "Tensid-Taschenbuch" (Handbook of Surfactants), Carl Hanser Verlag, Munich/Vienna 1981, and M. and J. Ash, "Encyclopedia of Surfactants", Vol I-III, Chemical Publishing Co., New York, 1980-81.

The agrochemical compositions will usually contain from 0.1 to 99 % by weight, preferably from 0.1 to 95 % by weight, of safener or mixture of safener and herbicide, from 1 to 99.9 % by weight, preferably from 5 to 99.8 % by weight, of a solid or liquid formulation assistant, and from 0 to 25 % by weight, preferably from 0.1 to 25 % by weight, of a surfactant.

Whereas it is preferred to formulate commercial products as concentrates, the end user will normally use dilute formulations.

The compositions may also contain further ingredients such as stabilisers, antifoams, viscosity regulators, binders, tackifiers, as well as fertilisers or other chemical agents.

Different methods and techniques may suitably be used for applying the safeners of formula II or compositions containing them for protecting cultivated plants from the harmful effects of herbicides of formula I, conveniently the following:

i) Seed dressing

a) Dressing the seeds with a wettable powder formulation of the compound of formula IIa, IIb or IIb₁ by shaking in a vessel until the safener is uniformly distributed on the surface of the seeds (dry treatment), using up to c. 1 to 500 g of compound of formula IIa, IIb or IIb₁

(4 g to 2 g of wettable powder) per 100 kg of seeds.

- b) Dressing seeds with an emulsifiable concentrate of the compound of formula IIa, IIb, IIb₁ by method a) (wet treatment).
- c) Dressing by immersing the seeds in a mixture containing 100-1000 ppm of compound of formula IIa, IIb or IIb₁ for 1 to 72 hours, leaving them wet or subsequently drying them (seed soaking).

Seed dressing or treatment of the germinated seedlings are naturally the preferred methods of application, as the safener treatment is fully concentrated on the target crop. Usually 1 to 1000 g, preferably 5 to 250 g, of safener is used per 100 kg of seeds. However, depending on the method employed, which also permits the use of other chemical agents or micronutrients, plus or minus deviations from the indicated limiting concentrations are possible (repeat dressing).

ii) Application as a tank mixture

A liquid formulation of a mixture of safener and herbicide (reciprocal ratio from 10:1 to 1:100) is used, the concentration of herbicide being from 0.005 to 5.0 kg/ha. This tank mixture is applied before or after sowing.

iii) Application in the furrow

The safener formulated as emulsifiable concentrate, wettable powder or granulate is applied to the open furrow in which the seeds have been sown. After covering the furrow, the herbicide is applied pre-emergence in conventional manner.

iv) Controlled release of safener

A solution of the compound of formula IIa, IIb or IIb₁ is applied to mineral granulate substrates or polymerised granulates (urea/formaldehyde) and allowed to dry. A coating may additionally be applied (coated granulates) which permits controlled release of the safener over a specific period of time.

The invention is illustrated in more detail by the following non-limitative Examples.

- 90 -

Formulation Examples for mixtures of formula I and safeners of formula IIa, IIb or IIb₁ (throughout, percentages are by weight)

F1. Emulsifiable concentrates	a)	b)	c)	d)
compound mixture	5 %	10%	25 %	50%
calcium dodecylbenzenesulfonate	6 %	8 %	6 %	8 %
polyethoxylated castor oil	4 %	_	4 %	4 %
(36 mol EO)				
octylphenol polyethoxylate	_	4 %	-	2 %
(7-8 mol EO)				
cyclohexanone	-	_	10 %	20 %
mixture of aromatic hydrocarbons	85 %	78 %	55 %	16%
C ₉ -C ₁₂				

Emulsions of any desired concentration can be prepared by diluting such concentrates with water.

F2. Solutions	a)	b)	c)	d)
compound mixture	5 %	10 %	50 %	90 %
1-methoxy-3-(3-methoxypropoxy)-	-	20 %	20 %	-
propane				
polyethylene glycol 400	20 %	10 %	-	-
N-methyl-2-pyrrolidone	-	_	30 %	10 %
mixture of aromatic hydrocarbons	75 %	60 %	-	-
C ₉ -C ₁₂				

The solutions are suitable for use as microdrops.

F3. Wettable powders	a)		b)		c)		d)	
compound mixture	5	8	25	&	50	윰	80	ક
sodium ligninsulfonate	4	용	-		3	용	-	
sodium laurylsulfate	2	ક	3	윰	-		4	용
sodium diisobutylnaphthalene								

- 91 -

sulfonate	-		6	용	5	윰	6 %	
octylphenol polyethoxylate	-		1	윰	2	용	-	
(7-8 mol EO)								
highly dispersed silica	1	ક	3	ક	5	용	10	용
kaolin	88	용	62	ક	35	ક	_	

The compound mixture is throughly mixed with the adjuvants and this mixture is ground in a suitable mill to give wettable powders which can be diluted with water to give suspensions of any desired concentration.

F4. Coated granulates	a)		b)		c)	
compound mixture	0.1	용	5	용	15	£
highly dispersed silica	0.9	용	2	윰	2	ક
inorganic carrier	99.0	ક	93	용	83	ક
(Ø 0.1-1 mm)						
e.g. CaCO ₃ or SiO ₂						

The compound mixture is dissolved in methylene chloride, the solution is sprayed on to the carrier, and the solvent is removed under vacuum.

F5. Coated granulates	a)		b)		c)	
compound mixture	0.1	용	5	윰	15	¥
polyethylene glycol 200	1.0	용	2	ક	3	용
highly dispersed silica	0.9	윰	1	윰	2	ક
inorganic carrier	98.0	ક	92	ક	80	용
(Ø 0.1 - 1 mm)						
e.g. CaCO ₃ or SiO ₂						

The finely ground compound mixture is uniformly applied in a mixer to the kaolin moistened with polyethylene glycol. Non-dusty coated granulates are obtained in this manner.

- 92 -

F6. Extruder granulates	a)		b)		c)		d)		
compound mixture	0.1	윰	3	ક	5	f	15	용	
sodium ligninsulfonate	1.5	ક	2	ક	3	용	4	ક્ર	
carboxymethyl cellulose	1.4	ક	2	ક	2	용	2	용	
kaolin	97.0	용	93	æ	90	용	79	8	

The compound mixture is mixed with the adjuvants and the mixture is moistened with water. This mixture is extruded and then dried in a stream of air.

F7. Dusts	a)	b)	c)
compound mixture	0.1	% 1	% 5 %
talcum	39.9	% 49	% 35 %
kaolin	60.0	% 50	% 60 %

Ready for use dusts are obtained by mixing the the active ingredient with the carriers on a suitable mill.

F8. Suspension concentrates	a)	b)	c)	d)
compound mixture	3 %	10 %	25 %	50 %
ethylene glycol	5 %	5 %	5 %	5 %
nonylphenol polyethoxylate	-	1 %	2 %	-
(15 mol EO)				
sodium ligninsulfonate	3 %	3 %	4 %	5 %
carboxymethyl cellulose	1 %	1 %	1 %	1 %
37% aqueous formaldehyde solution	0.2 %	0.2 %	0.2 %	0.2 %
silicone oil emulsion	0.8 %	0.8 %	0.8 %	0.8 %
water	87 %	79 %	62 %	38 %

The finely ground compound mixture is intimately mixed with the adjuvants to give a suspension concentrate from which suspensions of any desired concentration can be prepared by dilution with water.

- 93 -

The following Examples illustrate the ability of the safeners of formula IIa, IIb or IIb₁ to protect cultivated plants from the phytotoxic action of herbicides of formula L

Biological Examples

Examples B1 to B6: The test plants are raised under greenhouse conditions in plastic pots to the 2- to 3-leaf stage. The growth substrate is standard soil. To the test plants are applied at this stage on the one hand the herbicidal composition by itself and, on the other hand, the mixtures of the herbicidal composition with the substances to be tested as safeners. Application is made with an aqueous suspension of the test substances, prepared from a 25% wettable powder formulation (Example F3, b)) in 500 l of water/ha, with a standard spray nozzle. The rates of application are governed by the optimum rates determined according to field and greenhouse conditions. 3 weeks after application, the phytotoxic action of the herbicidal composition on the cultivated plants, e.g. maize and cereals, is evaluated on a percentage scale. 100 % = plant withered, 0 % = no phytotoxic action.

The results obtained in this test show that the damage caused by the herbicidal composition of formula I can be markedly reduced with the compounds of formula IIa, IIb and IIb₁. The following Examples demonstrate this effect in cereals and maize.

- 94 -

<u>Table B1:</u> Phytotoxic action of herbicide 1.010 singly and in admixture with safener 16.01 in cereals; postemergence application.

	Phytotoxic action:						
	herbicide	1.010	herbicide 1.010				
	AS [g/ha]		+			
			safener 1	6.01			
			AS [g/ha]			
	250	125	250	125			
			+	+			
test plants:			125	60			
winter barley	80	50	30	5			
summer wheat	60	30	0	0			
winter wheat	0	0	0	0			
Apera	90	80	95	90			
Alopecurus	80	40	50	30			
Avena	90	70	95	60			
Bromus	95	70	40	10			
Lolium	98	98	100	100			

<u>Table B2:</u> Phytotoxic action of herbicide 1.157, singly and in admixture with safener 16.01 in cereals; postemergence application.

	Phytotoxic action:						
	herbicide	1.157	herbicide 1.157				
	AS [g/ha]	+				
			safener 1	6.01			
			AS [g/ha]]			
	250	125	250	125			
			+	+			
test plants:			125	60			
winter barley	80	70	30	10			
summer wheat	70	70	40	30			

WO 96/11574		2199636	
	•	L 1 / / U J U	

Bromus

Lolium

- 95 -winter wheat Apera Alopecurus Avena

PCT/EP95/03935

Table B3: Phytotoxic action of herbicide 15.067, singly and in admixture with safener 16.01 in cereals; postemergence application.

	Phytotoxic action:						
	herbicide	15.067	herbicide 15.067				
	AS [g/ha]	l	•	+			
,			safener 1	6.01			
			AS [g/ha]				
	250	125	250	125			
			+	+			
test plants:			125	60			
winter barley	10	0	10	5			
summer wheat	30	. 15	0	0			
winter wheat	0	0	0	0			
Apera	90	80	80	60			
Alopecurus	60	30	30	10			
Avena	90	90	40	30			
Bromus	70	70	0	0			
Lolium	98	95	95	90			

- 96 -

<u>Table B4:</u> Phytotoxic action of herbicide 1.010, singly and in admixture with safener 16.01 in maize; postemergence application.

	Phytotoxi herbicide AS [g/ha]	1.010	herbicide 1.010			
	250	105	safener 1 AS [g/ha	1		
	250	125	250 +	125 +		
test plants:			250	125		
maize	75	70	30	20		
Digitaria	100	95	100	98		
Echinochloa	100	90	100	90		
Panicum	100	100	100	100		
Setaria	90	70	90	75		

<u>Table B5:</u> Phytotoxic action of herbicide 1.157, singly and in admixture with safener 16.01 in maize; postemergence application.

	Phytotox	ic action:			
	herbicide	1.157	herbicide 1.157		
	AS [g/ha]]	•	+	
			safener 1	6.01	
			AS [g/ha]		
	125	60	125	60	
			+	+	
test plants:			125	60	
maize	95	90	45	10	
Digitaria	100	90	100	90	
Echinochloa	100	100	100	100	

- 97 -

Panicum	100	100	100	100
Setaria	75	40	30	20

<u>Table B6:</u> Phytotoxic action of herbicide 15.067, singly and in admixture with safener 16.01 in maize; postemergence application.

	Phytotoxic	action:			
	herbicide 15.067 AS [g/ha]		herbicide 15.	herbicide 15.067	
			+ safener 16.01		
	safener 16.01 AS [g/ha]				
	125	60	125	60	
			+	+	
test plants:			125	60	
maize	75	35	10	0	
Digitaria	95	90	90	60	
Echinochloa	90	70	90	70	
Panicum	100	90	100	80	
Setaria	80	50	85	40	

The same results are obtained by formulating the mixtures in accordance with Examples F1, F2 and F4 to F8.

THIS PAGE BLANK (US. 073)

What is claimed is:

- 1. A selective herbicidal composition comprising, in addition to customary inert formulation assistants, a mixture of
- a) a herbicidally effective amount of a herbicide of formula I

wherein

$$R_1$$
 is the $(R_4)_n$ or $(R_4)_n$ group;

the substituents R_4 are each independently of one another halogen, nitro, cyano, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_{10} alkoxy, C_1 - C_4 haloalkoxy, C_3 - C_6 alkenyloxy, C_1 - C_4 alkoxy- C_2 - C_4 alkoxy, C_3 - C_6 alkynyloxy, C_1 - C_4 alkylcarbonyl, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, amino, C_1 - C_4 alkylamino or di- C_1 - C_4 alkylamino;

$$R_5$$
 is the $-x_1$ or $-x_1$ (R_6) q group;

n is 0, 1, 2, 3 or 4;

m is 0 or 1, the sum of m and n being 0, 1, 2, 3 or 4;

q is 0, 1, 2 or 3;

 X_1 is oxygen, sulfur, -CH₂- or -N(R₇)-;

the substituents R_6 are each independently of one another C_1 - C_4 alkyl, halogen, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, nitro, cyano, C_1 - C_4 alkoxycarbonyl, amino, C_1 - C_4 alkylamino or di- C_1 - C_4 alkylamino;

R₇ is hydrogen, C₁-C₄alkyl, formyl or C₁-C₄alkylcarbonyl;

A and B are each independently of the other hydrogen, alkyl, alkenyl, alkynyl, alkoxyalkyl, alkylthioalkyl or cycloalkyl, or substituted or unsubstituted aryl; or A and B, taken together, form the divalent radical of a saturated or unsaturated and unsubstituted or substituted mono-, bi-, tri- or polycyclic system;

THIS PAGE BLANK (USOTO)

- 99 -

G is hydrogen or a group -CO-R₁₈ (a),
$$M-R_{19}$$
 (b), -SO₂-R₂₀ (c), R_{22} (d),

$$C-N$$
 R_{24} (e) or X (f);

L and M are each independently of the other oxygen or sulfur;

R₁₈ is halogen-substituted alkyl, alkenyl, alkoxyalkyl, alkylthioalkyl, polyalkoxyalkyl or cycloalkyl, which may contain hetero atoms; unsubstituted or substituted phenyl, unsubstituted or substituted phenylalkyl, substituted heteroaryl, substituted phenoxyalkyl, or substituted heteroaryloxyalkyl;

R₁₉ is halogen-substituted alkyl, alkenyl, alkoxyalkyl or polyalkoxyalkyl, or unsubstituted or substituted phenyl or benzyl;

 R_{20} , R_{21} and R_{22} are each independently of one another unsubstituted or halogen-substituted alkyl, alkoxy, alkylamino, dialkylamino, alkylthio, alkenylthio, alkynylthio or cycloalkylthio, or unsubstituted or substituted phenyl, phenoxy or phenylthio;

 R_{23} and R_{24} are each independently of the other hydrogen, unsubstituted or halogen-substituted alkyl, alkenyl, alkoxy or alkoxyalkyl, unsubstituted or substituted phenyl or benzyl; or

R₂₃ and R₂₄, taken together, form an alkenyl radical, which may contain oxygen as hetero atom; and

X is a metal ion equivalent or an ammonium ion; as well as salts and diastereoisomers of the compounds of formula I; and

b) to antagonise the herbicide, an antidotally effective amount of either a quinoline derivative of formula IIa

- 100 -

wherein

 R_{10} is hydrogen, C_1 - C_8 alkyl, or C_1 - C_8 alkyl which is substituted by C_1 - C_6 alkoxy or C_3 - C_6 alkenyloxy; and X_2 is hydrogen or chloro; of a 1-phenylazole-3-carboxylic acid derivative of formula IIb

$$R_{11}$$
 R_{12}
 R_{13}
 R_{13}
 R_{14}
 R_{15}
 R_{15}

wherein

E is nitrogen or methine;

 R_{11} is -CCl₃ or unsubstituted or halogen-substituted phenyl; R_{12} and R_{13} are each independently of the other hydrogen or halogen; and R_{14} is C_1 -C₄alkyl; or of a compound of formula IIb₁

$$R_{67}$$
COOR₆₆
 R_{68}
 R_{12}
 R_{13}
(IIb₁),

wherein R_{12} and R_{13} have the meanings given above, and R_{66} , R_{67} and R_{68} are each independently of one another C_1 - C_4 alkyl.

2. A composition according to claim 1, wherein R_1 is the $\binom{(R_4)_n}{(R_5)_n}$ group.

- 3. A composition according to claim 2, wherein R_4 is C_1 - C_4 alkyl; n is 1, 2, 3 or 4; and m is 0.
- 4. A composition according to claim 2, wherein R_5 is the X_1 X_1 X_2 X_3 X_4 X_5 X_5
- 5. A composition according to claim 2, wherein R_4 is C_1 - C_4 alkyl, trifluoromethyl, fluoro, chloro, bromo, C_1 - C_{10} alkoxy, allyloxy, propargyloxy, difluoromethoxy, trifluoromethoxy, methoxyethylenoxy, cyano, nitro, di- C_1 alkylamino or di- C_2 alkylamino, acetyl, C_1 - C_3 alkoxycarbonyl, methylsulfonyl, methylsulfinyl or methylmercapto; R_5 is the X_1 - X_2 - X_3 - X_4 - X_4 - X_5 - $X_$

or -N(CHO)-; R₆ is fluoro, chloro or trifluoromethyl; n is 0, 1, 2 or 3; m is 0 or 1; and o is 0, 1 or 2, the sum of m and n being 0, 1, 2 or 3.

6. A composition according to claim 1, wherein R_1 is the $(R_4)_n$ group; R_4 is fluoro, chloro, methyl, nitro, trifluoromethyl or methylsulfonyl; n is 0, 1 or 2; and m is 0.

- 7. A composition according to claim 1, wherein R_1 is the is methyl, methoxy, fluoro, chloro or nitro; and n is 0 or 1.
- 8. A composition according to claim 1, wherein the safener is a compound of formula IIc

9. A composition according to claim 1, wherein the safener is a compound of formula IId

2199636

- 102 -

$$\begin{array}{c} \text{CI} \\ \\ \text{O-CH}_2\text{-C(O)-O-CH(CH}_3\text{)-CH}_2\text{-O-CH}_2\text{CH=-CH}_2 \end{array}$$

10. A composition according to claim 1, wherein the safener is a compound of formula IIe

- 11. A method of selectively controlling weeds and grasses in crops of cultivated plants, the seeds or the locus thereof, concurrently or separately, with an effective amount of a herbicide of formula I as claimed in claim 1 and, to antagonise said herbicide, an antidotally effective amount of a safener of formula IIa, IIb or IIb₁ as claimed in claim 1.
- 12. A method according to claim 11, which comprises treating crops of cultivated plants or areas intended for cropping with cultivated plants, with 0.001 to 2 kg/ha of a herbicide of formula I and an amount of 0.001 to 0,5 kg/ha of a safener of formula IIa, IIb or IIb₁.
- 13. A method according to claim 11, wherein the cultivated plants are maize and cereals.